

INTRODUCTION TO THE LOCAL LANGLANDS CORRESPONDENCE

MASAO OI

CONTENTS

1. Week 1: Course overview	3
1.1. Class field theory	3
1.2. What is the Langlands correspondence?	5
1.3. Local-global principle in number theory	6
1.4. What is the local Langlands correspondence?	7
2. Week 2: Overview of local class field theory	9
2.1. Local fields and CDVR	9
2.2. Extension of local fields	10
2.3. Galois groups and Weil groups of local fields	13
2.4. Local class field theory	14
3. Week 3: Representations of locally profinite groups	16
3.1. Locally profinite groups	16
3.2. Smooth representations of locally profinite groups	17
3.3. Frobenius reciprocity	19
3.4. Representations of profinite groups	19
3.5. Contragredient representation	20
3.6. Irreducible representations and Schur's lemma	21
4. Week 4: Irreducible smooth representations of $\mathrm{GL}_2(F)$	23
4.1. Recap on irreducible representations of $\mathrm{GL}_2(\mathbb{F}_q)$	23
4.2. Principal series representations of $\mathrm{GL}_2(F)$	24
4.3. Depth-zero supercuspidal representations of $\mathrm{GL}_2(F)$	25
4.4. Depth of representations	26
4.5. Simple supercuspidal representations	27
5. Week 5: Representation of Weil groups	29
5.1. Representations absolute Galois groups	29
5.2. Galois group vs. Weil group	30
5.3. More about Weil groups	31
5.4. Grothendieck's monodromy theorem	32
5.5. Weil–Deligne representations	34
6. Week 6: Local Langlands correspondence for GL_n	36
6.1. Local Langlands correspondence for GL_n	36
6.2. Example: the case of GL_2	37
6.3. Idea of the characterization of LLC for GL_n	38
6.4. Local L -factors and ε -factors	39
6.5. Local L -factors and ε -factors for pairs	41

1. WEEK 1: COURSE OVERVIEW

1.1. Class field theory. Let us begin with the following very famous and classical theorem in elementary number theory.

Theorem 1.1. *The number of the solutions to the equation $x^2 - 2 = 0$ in \mathbb{F}_p is given as follows:*

$$|\{x \in \mathbb{F}_p \mid x^2 - 2 = 0\}| = \begin{cases} 2 & \text{if } p \equiv 1, 7 \pmod{8}, \\ 0 & \text{if } p \equiv 3, 5 \pmod{8}, \\ 1 & \text{if } p = 2. \end{cases}$$

This theorem is called *the second supplement to the quadratic reciprocity law* (see, e.g., [Ser73, Chapter I, §3]). In fact, more generally, the general quadratic reciprocity law implies the following:

Theorem 1.2. *Let $a \in \mathbb{Z}$ be an integer. Then there exists a positive integer $N \in \mathbb{Z}_{>0}$ such that the number $|\{x \in \mathbb{F}_p \mid x^2 - a = 0\}|$ depends only on the modulo N of p .*

For example, Theorem 1.1 says that N can be taken to be 8 when $a = 2$.

Exercise 1.3. (1) Explain the statement of the quadratic reciprocity law.

(2) Determine the number N in Theorem 1.2 using the quadratic reciprocity law.

Next let us consider the equation $x^3 - 2 = 0$. Can we find a simple description of the numbers of the solutions to this equation in \mathbb{F}_p like above? In fact, the answer is NO! More precisely, there does not exist a positive integer $N \in \mathbb{Z}_{>0}$ such that the number $|\{x \in \mathbb{F}_p \mid x^3 - 2 = 0\}|$ depends only on the modulo N of p .

What causes such a difference between the quadratic and the cubic cases? To explain it, let us think about how to prove Theorem 1.1 from a modern viewpoint based on algebraic number theory. (In the following, we appeal to some basics of algebraic number theory. But it's not a material necessary for this course. If you are not familiar with them, please try to feel just its flavor.)

Since the equality $|\{x \in \mathbb{F}_2 \mid x^2 - 2 = 0\}| = 1$ is obvious, let us suppose that p is an odd prime number. Then Theorem 1.1 is rephrased as follows:

$$\mathbb{F}_p \text{ has a square root of } 2 \text{ if and only if } p \equiv \pm 1 \pmod{8}.$$

Noting this, let us introduce the quadratic extension $K := \mathbb{Q}(\sqrt{2})$ of \mathbb{Q} obtained by adding a square root $\sqrt{2}$ of 2. The ring of integer \mathcal{O}_K in K is given by $\mathbb{Z}[\sqrt{2}]$. Because the quadratic extension K/\mathbb{Q} is unramified outside 2, any odd prime number p has only the following two possibilities about the ideal $p\mathcal{O}_K$ of \mathcal{O}_K generated by p :

- $p\mathcal{O}_K$ is a prime (maximal) ideal of \mathcal{O}_K (p “inerts” in K), or
- $p\mathcal{O}_K$ is the product $\mathfrak{p}_1\mathfrak{p}_2$ of two different prime ideals \mathfrak{p}_1 and \mathfrak{p}_2 of \mathcal{O}_K (p “splits completely” in K).

Let us look at the quotient ring $\mathcal{O}_K/p\mathcal{O}_K$. This ring is

- a field if p inerts in K , and
- the product of two fields ($\mathcal{O}_K/\mathfrak{p}_1$ and $\mathcal{O}_K/\mathfrak{p}_2$) if p splits completely in K .

On the other hand,

$$\begin{aligned} \mathcal{O}_K/p\mathcal{O}_K &= \mathbb{Z}[\sqrt{2}]/p\mathbb{Z}[\sqrt{2}] \cong (\mathbb{Z}[x]/(x^2 - 2))/p(\mathbb{Z}[x]/(x^2 - 2)) \\ &\cong \mathbb{F}_p[x]/(x^2 - 2). \end{aligned}$$

The right-hand side is

- a field (a quadratic extension of \mathbb{F}_p) if \mathbb{F}_p does not have a square root of 2, and
- the product of two fields (both \mathbb{F}_p) if \mathbb{F}_p has a square root of 2.

Hence, in summary, we see that

\mathbb{F}_p has a square root of 2 if and only if p splits completely in K .

Recall that each odd prime number p gives rise to a special element Frob_p of $\text{Gal}(K/\mathbb{Q})$, called *Frobenius element* (again note that K/\mathbb{Q} is unramified outside 2). The important property of the Frobenius is that it knows whether p splits completely or not. More precisely,

p splits completely in K if $\text{Frob}_p = \text{id}$.

So, our task is now reduced to investigate when $\text{Frob}_p = \text{id}$.

In fact, the argument so far can be carried out in general (e.g., for $x^3 - 2 = 0$ by replacing K with the smallest factorization field of $x^3 - 2 = 0$) more or less. But here we reach the stage where a special nature of the equation $x^2 - 2 = 0$ comes into play. The point is that the quadratic extension K/\mathbb{Q} is abelian, i.e., its Galois group $\text{Gal}(K/\mathbb{Q})$ is abelian. In general, by the Kronecker–Weber theorem, any abelian extension of \mathbb{Q} is contained in a cyclotomic field $\mathbb{Q}(\mu_N)$ (μ_N denotes the set of N -th roots of unity). The Galois group of $\mathbb{Q}(\mu_N)/\mathbb{Q}$ is given by $(\mathbb{Z}/N\mathbb{Z})^\times$; by choosing a primitive N -th root ζ_N of unity, it is described as follows:

$$\text{Gal}(\mathbb{Q}(\mu_N)/\mathbb{Q}) \cong (\mathbb{Z}/N\mathbb{Z})^\times : [\zeta_N \mapsto \zeta_N^i] \mapsto i.$$

Under this identification, the Frobenius element Frob_p on the left-hand side is mapped to $p \in (\mathbb{Z}/N\mathbb{Z})^\times$ on the right-hand side (as long as p is unramified, which is equivalent to that p does not divide N).

In our situation, actually we have $\mathbb{Q}(\sqrt{-2}) \subset \mathbb{Q}(\mu_8)$. More precisely, under the Galois theory, $\mathbb{Q}(\sqrt{-2})$ is the subfield of $\mathbb{Q}(\mu_8)$ corresponding to the subgroup $\{\pm 1\}$ of $\text{Gal}(\mathbb{Q}(\mu_8)/\mathbb{Q}) \cong (\mathbb{Z}/8\mathbb{Z})^\times$. Hence the Galois group $\text{Gal}(K/\mathbb{Q})$ is identified with the quotient of $(\mathbb{Z}/8\mathbb{Z})^\times$ by $\{\pm 1\}$. Thus we conclude that

$$\text{Frob}_p = \text{id} \text{ if and only if } p \equiv \pm 1 \pmod{8}.$$

Hence this completes the proof of Theorem 1.1.

The classical class field theory enables us to do a similar thing for more general number fields (finite extensions of \mathbb{Q}).

Theorem 1.4 (class field theory). *Let F be a number field. Let F^{ab} be the maximal abelian extension of F . Then there exists a natural surjective continuous homomorphism*

$$\text{Art}_F: \mathbb{A}_F^\times \rightarrow \text{Gal}(F^{\text{ab}}/F),$$

which kernel is explicitly described.

Here, I do not explain the meaning of “natural” (it is formulated as the compatibility with the local class field theory, which will be explained later) nor even what “ \mathbb{A}_F ” on the source of the map is. But I just want to emphasize that this “ \mathbb{A}_F ” (which is called the adèle ring of F) is defined only using the intrinsic data of the original object F . So, class field theory describes how the field F extends to a larger abelian field only by appealing to the internal data of F , which is much easier to grasp. For example, when $F = \mathbb{Q}$, the map Art_F exactly gives rise to the above-mentioned isomorphism $(\mathbb{Z}/N\mathbb{Z})^\times \cong \text{Gal}(\mathbb{Q}(\mu_N)/\mathbb{Q})$ by taking an appropriate finite quotient.

If we try to imitate the above discussion in the case of the equation $x^3 - 2 = 0$, we immediately notice that the last part does not work because the smallest splitting field $\mathbb{Q}(\sqrt[3]{2}, \mu_3)$ of the equation $x^3 - 2 = 0$ is not abelian over \mathbb{Q} ; its Galois group is given by \mathfrak{S}_3 .

1.2. What is the Langlands correspondence? Then, is it impossible to find any beautiful law on the behavior of the number $|\{x \in \mathbb{F}_p \mid x^3 - 2 = 0\}|$ over prime numbers p ? In fact, the following holds:

Theorem 1.5. *We let $\sum_{n=1}^{\infty} a_n q^n$ be the infinite series given by the following infinite product:*

$$q \cdot \prod_{n=1}^{\infty} (1 - q^{6n}) \cdot (1 - q^{18n}) = \sum_{n=1}^{\infty} a_n q^n.$$

Then, for any prime number $p \neq 2, 3$, we have

$$|\{x \in \mathbb{F}_p \mid x^3 - 2 = 0\}| = 1 + a_p.$$

(See, e.g., [DS05, Section 4.11] for the more general case of $x^3 - a = 0$.)

Let us also introduce a different, but similar, phenomenon. We consider the following equation:

$$E: y^2 + y = x^3 - x^2.$$

The set of solutions of this equation forms a curve, which is called an *elliptic curve*. Let us think about the solutions in \mathbb{F}_p :

$$E(\mathbb{F}_p) := \{(x, y) \in \mathbb{F}_p \times \mathbb{F}_p \mid y^2 + y = x^3 - x^2\}.$$

Note that, in this case, the equation is not one-variable. So we do not even have a simple interpretation of the set $E(\mathbb{F}_p)$ in terms of field extensions of \mathbb{Q} . (In the case of $x^3 - 2 = 0$, although we cannot apply the class field theory, we can still relate the number $|\{x \in \mathbb{F}_p \mid x^3 - 2 = 0\}|$ to how p decomposes into prime ideals in the smallest splitting field of $x^3 - 2 = 0$.) Nevertheless, we have the following:

Theorem 1.6. *We let $\sum_{n=1}^{\infty} a_n q^n$ be the infinite series given by the following infinite product:*

$$\sum_{n=1}^{\infty} a_n q^n = q \cdot \prod_{n=1}^{\infty} (1 - q^n)^2 \cdot (1 - q^{11n})^2.$$

Then, for any prime number $p \neq 11$, we have

$$|E(\mathbb{F}_p)| = 1 + p - a_p.$$

In Theorems 1.5 and 1.6, by putting $q := \exp(2\pi iz)$ (for $z \in \mathbb{C}$), we may regard the infinite series as functions on the complex upper-half plane. In fact, they are examples of so-called “modular forms”, which is a holomorphic function on the complex upper-half plane equipped with a lot of symmetry. Both elliptic curves and modular forms have been investigated in the context of number theory for a long time. A priori, they are totally different objects; elliptic curves are purely-algebraic while modular forms are purely-analytic, at least from the above descriptions. However, they are actually related in a surprising way as above.

All the phenomena introduced so far (Theorems 1.1, 1.5, 1.6) can be thought of as special cases of the *Langlands correspondence*. The Langlands correspondence is a vast, but conjectural, framework which connects two completely different mathematical objects: on the one hand are *automorphic representations* and on the other hand are *Galois representations*:

$$(\text{automorphic representations}) \quad \overset{\text{Langlands correspondence}}{\longleftrightarrow} \quad (\text{Galois representations})$$

Roughly speaking, an automorphic representation is an irreducible representation of $\mathrm{GL}_n(\mathbb{A}_F)$ realized in the space of automorphic forms, which are generalization of modular forms, and a Galois representation is an n -dimensional continuous¹ representation of the absolute Galois group $\mathrm{Gal}(\overline{F}/F)$.

The important viewpoint here is not to look at the Galois group itself, but to consider representations of the Galois group. Recall that representation theory is a very strong tool (or even a modern “formulation”) for studying non-abelian groups. For example, when $n = 1$, we have $\mathrm{GL}_1(\mathbb{A}_F) = \mathbb{A}_F^\times$; this implies an automorphic representation of $\mathrm{GL}_1(\mathbb{A}_F)$ is just a character of \mathbb{A}_F^\times . On the other hand, when the dimension of a Galois representation is 1, it must be a character, hence it necessarily factors through the maximal abelian quotient of $\mathrm{Gal}(\overline{F}/F)$, i.e., $\mathrm{Gal}(\overline{F}^{\mathrm{ab}}/F)$. Thus the Langlands correspondence in this case says that the characters of \mathbb{A}_F^\times and $\mathrm{Gal}(\overline{F}^{\mathrm{ab}}/F)$ nicely correspond. This is exactly implied by the isomorphism $\mathbb{A}_F^\times \cong \mathrm{Gal}(\overline{F}^{\mathrm{ab}}/F)$ of class field theory.

When $n = 2$, the Shimura–Taniyama conjecture, which plays a crucial role in the proof of Fermat’s conjecture, is also regarded as a special case of the Langlands correspondence. Theorem 1.6 is an example of the Shimura–Taniyama conjecture.

Other than these examples, It is known that various phenomena in number theory can be explained in a sophisticated way by appealing to the prediction of the Langlands correspondence. Therefore, one of the most important objectives in the modern number theory is to establish the Langlands correspondence.

Exercise 1.7. By looking at “LMFDB” (which is an online database of modular forms, elliptic curves, and so on), we can find a lot of examples of elliptic curves and modular forms which “correspond”. For example, the elliptic curve and the modular form considered in Theorem 1.6 are labelled by “11.a3” and “11.2.a.a”, respectively. I just randomly chose the following elliptic curve from this database: $y^2 + xy + y = x^3 - x$. Try to find the modular form corresponding to this elliptic curve using LMFDB (please explain how you arrive at it).

1.3. Local-global principle in number theory. Then, what is the “local” Langlands correspondence in the course title? To explain this, let us briefly talk about the philosophy of the local-global principle in number theory. Recall that the real number field \mathbb{R} is the completion of the rational number field \mathbb{Q} with respect to the normal metric on \mathbb{Q} . We note that \mathbb{R} is not the only field obtained by such a procedure from \mathbb{Q} . Indeed, \mathbb{Q} possesses non-trivial metrics other than the normal metric. For each fixed prime number p , if we put $|p^r \cdot \frac{n}{m}|_p := p^{-r}$ (here, n and m are integers prime to p), then $|\cdot|_p$ gives a well-defined metric on \mathbb{Q} called the p -adic metric. If we complete \mathbb{Q} with respect to the p -adic metric, we obtain a locally compact field different to \mathbb{R} , which is called the p -adic number field and denoted by \mathbb{Q}_p . The fundamental philosophy in number theory is that any problem on the rational number field \mathbb{Q} should be able to be understood through its analog for \mathbb{R} and \mathbb{Q}_p for all prime numbers p ; this is the idea of “local-global” in number theory.

problem on \mathbb{Q} $\xleftrightarrow{\text{local-global principle}}$ problems on \mathbb{R} and \mathbb{Q}_p (for all p)

For example, the local analog of the class field theory is the *local class field theory*, which says that, for any p -adic field F (i.e., a finite extension of \mathbb{Q}_p), we have a natural injective

¹It is very important which kind of coefficient field/topology we adopt when we consider a representation of $\mathrm{Gal}(\overline{F}/F)$. But let us just ignore this subtlety here.

homomorphism

$$\text{Art}_F: F^\times \rightarrow \text{Gal}(F^{\text{ab}}/F)$$

with dense image.

Both automorphic representations and Galois representations are objects related to the rational number field \mathbb{Q} (or, more generally, any number field F). Thus it is natural to think about the analog of the Langlands correspondence for \mathbb{R} or \mathbb{Q}_p (or, more generally, any local field of characteristic zero, which means a finite extension of \mathbb{R} or \mathbb{Q}_p); this is what is called the *local Langlands correspondence (LLC)*. This also generalized the local class field theory.

1.4. What is the local Langlands correspondence? Let us explain the LLC a bit more precisely. In the following, we let F be any p -adic field, i.e., a finite extension of \mathbb{Q}_p . The LLC is a natural correspondence between the set of “irreducible admissible representations” of $\text{GL}_n(F)$ and the set of “ n -dimensional Weil–Deligne representations”:

$$(\text{irred. adm. reps. of } \text{GL}_n(F)) \xleftrightarrow{\text{LLC}} (n\text{-dim. WD reps.})$$

Here, roughly speaking,

- an *irreducible admissible representation* of $\text{GL}_n(F)$ means an irreducible representation of the group $\text{GL}_n(F)$ on a \mathbb{C} -vector space equipped with a certain finiteness condition (this can be thought of as the local version of an automorphic representation);
- a *Weil–Deligne representation* is a modified version of the notion of a continuous representation of $\text{Gal}(\overline{F}/F)$.

Now recall that the starting point of our discussion was how to understand the absolute Galois group $\text{Gal}(\overline{F}/F)$. The point of class field theory is that it can be understood through a much easier object F^\times . However, at this point, we notice the following:

- The automorphic side of LLC is not so obvious to understand as in the case of F^\times . So we may also think that LLC enables us to investigate irreducible admissible representations of $\text{GL}_n(F)$ through the Galois side, which consists of arithmetic objects.
- The automorphic side of LLC makes sense even if we replace GL_n with more general groups.

Keeping these observations in mind, let us present a naive formulation of LLC in general:

Conjecture 1.8 (local Langlands conjecture, naive form). *Let G be a reductive group defined over F . Then there exists a natural map from the set of irreducible admissible representations of $G(F)$ to the set of “ L -parameters” of G .*

For general G , we can no longer say that one of the automorphic or Galois sides is particularly easier than the other side. Therefore the local Langlands correspondence is very important not only from number-theoretic viewpoint, but also representation-theoretic viewpoint (representation theory of p -adic reductive groups).

At present, LLC is still conjectural in general, but has been constructed for several specific groups. For example,

- GL_n by Harris–Taylor [HT01], Henniart [Hen00],
- SO_n and Sp_{2n} (quasi-split) by Arthur [Art13],
- U_n (quasi-split) by Mok [Mok15],
- and so on...

On the other hand, there are also approaches for specific classes of irreducible admissible representations. For example,

- the classical construction by Satake for unramified representations,
- regular depth-zero supercuspidal representations by DeBacker–Reeder [DR09],
- regular (positive-depth) supercuspidal representations by Kaletha [Kal19],
- and so on...

The aims of this course to understand the following:

- A naive formulation of LLC in general. For this, I will explain some basics of representation theory of p -adic reductive groups (such as the notion of admissible representations) and also representations theory of local Galois groups (especially, Weil–Deligne representations etc).
- The precise formulation (characterization) of LLC for GL_n given by [HT01] and [Hen00]. For this, I will explain more details of representaiton theory of p -adic reductive groups by focusing on the case of GL_n (so-called “Bernstein–Zelevinsky classification”). It is far beyond my ability to explain the construction of LLC, so I’m not going to touch it.
- The precise formulation (characterization) of LLC for quasi-split classical groups given by [Art13] and [Mok15]. For this, I will explain basics about harmonic analysis on p -adic reductive groups including the Harish–Chandra characters of representaitons etc.
- Recent developments on explicit construction of LLC for certain supercuspidal representations by [DR09], [Kal19], etc.

Of course, this plan must be too ambitious. Let’s see how much I can achieve...

2. WEEK 2: OVERVIEW OF LOCAL CLASS FIELD THEORY

2.1. Local fields and CDVR. We briefly review some basic facts about local fields (see, e.g., [Ser79, Chapters 1, 2] or [Wei74, Chapter I]).

We first introduce the *p-adic number field* \mathbb{Q}_p . Recall that the real number field \mathbb{R} is the completion of the rational number field \mathbb{Q} with respect to the normal metric on \mathbb{Q} . In fact, there is a different way of completing \mathbb{Q} ; for each prime number p , we put

$$|p^r \cdot \frac{n}{m}|_p := p^{-r}$$

(here, n and m are integers prime to p). Then $|\cdot|_p$ gives a well-defined metric on \mathbb{Q} called the *p-adic metric*. If we complete \mathbb{Q} with respect to the *p-adic metric*, we obtain a locally compact field different to \mathbb{R} , which is called the *p-adic number field* and denoted by \mathbb{Q}_p .

Local fields are generalizations of these fields.

Definition 2.1 (local field). We say that a field F is a *local field* if it is a nondiscrete locally compact topological field.

Fact 2.2. Any local field is isomorphic to one of the following:

- \mathbb{R} or \mathbb{C} (archimedean);
- a finite extension of \mathbb{Q}_p (nonarchimedean, characteristic 0);
- a finite extension of $\mathbb{F}_p((t))$ (nonarchimedean, characteristic p).

One notable characterization of a local field is that it is the completion of a *global field* (i.e., a finite extension of \mathbb{Q} or $\mathbb{F}_p(t)$) with respect to a nontrivial metric. Thus, from the viewpoint of “global” number theory, both archimedean and nonarchimedean local fields have equal importance. However, in this course, we focus only on nonarchimedean local fields (and often assume even that characteristic is zero).

Let us introduce more ring-theoretic description of nonarchimedean local fields.

Definition 2.3 (DVR (discrete valuation ring)). Let F be a field. We say that a group homomorphism $v: F^\times \rightarrow \mathbb{Z}^\times$ is a *discrete valuation* of F if it is surjective and satisfies the following condition:

$$\text{For any } x, y \in F, \text{ we have } v(x + y) \geq \min\{v(x), v(y)\},$$

where we put $v(0) := \infty$. When F is equipped with a discrete valuation v , the set

$$\{x \in F \mid v(x) \geq 0\}$$

forms a subring of F , called the *valuation ring* F (with respect to v). If a ring \mathcal{O} is obtained as the valuation ring of a field with respect to its discrete valuation, we call it a *discrete valuation ring (DVR)*.

Fact 2.4. Let \mathcal{O} be a ring. Then \mathcal{O} is a DVR if and only if it is a PID with unique nonzero prime (hence maximal) ideal.

When \mathcal{O} is a DVR with discrete valuation v , its subset

$$\{x \in F \mid v(x) = 0\}$$

forms the multiplicative group of units \mathcal{O}^\times . The maximal ideal of \mathcal{O} is given by

$$\mathfrak{p} = \{x \in F \mid v(x) \geq 1\}.$$

Any generator of the maximal ideal \mathfrak{p} is often referred to as a *uniformizer* of \mathfrak{p} . If we fix a uniformizer ϖ of \mathfrak{p} , then any nonzero ideal of \mathcal{O} is expressed as ²

$$\mathfrak{p}^n = \{x \in F \mid v(x) \geq n\} = \varpi^n \mathcal{O}.$$

We call \mathcal{O}/\mathfrak{p} the *residue field* of \mathcal{O} .

Now let F be a fractional field of a DVR \mathcal{O} with discrete valuation v . Then we can equip F with a metric $|x| := r^{v(x)}$ ($|0| := 0$) by choosing any real number $r \in (0, 1)$. If we let \hat{F} be the completion of F with respect to this metric, \hat{F} naturally has a structure of a topological field. Moreover, we can equip \hat{F} with a discrete valuation which extends v ; the valuation ring of \hat{F} is given by the closure of \mathcal{O} in \hat{F} . By noting that $\{\mathfrak{p}^n\}_{n \in \mathbb{Z}_{\geq 0}}$ forms a fundamental system of open neighborhoods of 0 in \mathcal{O} , we can see that the closure of \mathcal{O} in \hat{F} is nothing but

$$\hat{\mathcal{O}} := \varprojlim_n \mathcal{O}/\mathfrak{p}^n,$$

where the transition map $\mathcal{O}/\mathfrak{p}^{n+1} \rightarrow \mathcal{O}/\mathfrak{p}^n$ is given by the natural surjection.

We say that a DVR \mathcal{O} is *complete* (and say \mathcal{O} is a *CDVR*) if $\hat{\mathcal{O}} = \mathcal{O}$.

Fact 2.5. *Let F be a field. Then F is a nonarchimedean local field if and only if F is a fractional field of CDVR (“CDVF”) with finite residue field.*

Remark 2.6. When F is a nonarchimedean local field with valuation ring \mathcal{O} and maximal ideal \mathfrak{p} , the characteristics of $(F, \mathcal{O}/\mathfrak{p})$ must be either $(0, p)$ (called *mixed characteristic*) or (p, p) (called *equal characteristic*). According to a classification result mentioned above, F is mixed characteristic if and only if it is a finite extension of \mathbb{Q}_p . In this case, we often say that F is a *p-adic field* (but this terminology depends on people).

Let F be a nonarchimedean local field. Recall that the absolute Galois group of F is, by definition, the Galois group $\Gamma_F := \text{Gal}(F^{\text{sep}}/F)$ of the separable closure F^{sep} of F . ³ The separable closure F^{sep} is given by the direct limit (union) of all finite separable (Galois) extensions of F . We define F^{ab} to be the *maximal abelian extension* of F in F^{sep} , i.e., the direct limit (union) of all finite abelian extensions of F . (Note that this makes sense since the composite field of any two finite abelian extensions is again a finite abelian extension.) Then the Galois group $\text{Gal}(F^{\text{ab}}/F)$ is identified with the maximal abelian quotient of Γ_F , i.e., $\Gamma_F/[\Gamma_F, \Gamma_F]$.

2.2. Extension of local fields.

Fact 2.7. *Let \mathcal{O}_F be a CDVR with fractional field F . Let E/F be a finite separable extension of rank n . Then the integral closure of \mathcal{O}_F in E (write \mathcal{O}_E) is a CDVR. Moreover, \mathcal{O}_E is a free \mathcal{O}_F -module of rank $[E : F]$.*

By this fact, it makes sense to refer to \mathcal{O}_F as the *ring of integers in F* .

Let E/F be a finite separable extension of non-archimedean local fields of degree n . Let \mathcal{O}_F be the ring of integers in F , \mathfrak{p}_F the maximal ideal of \mathcal{O}_F , $k_F := \mathcal{O}_F/\mathfrak{p}_F$ the residue field. Also for E , we define \mathcal{O}_E , \mathfrak{p}_E , and k_E in a similar way. We introduce two invariants of the extension E/F :

²Another popular symbol for a uniformizer is π , but we often use ϖ in our area (representation theory of p -adic groups) in order to reserve π to denote a representation.

³Another standard symbol for the absolute Galois group is “ G_F ”, but we avoid it because we want to use “ G ” for a reductive group over F .

- The ideal $\mathfrak{p}_F \mathcal{O}_E$ of \mathcal{O}_E is of the form \mathfrak{p}_E^e , where $e \in \mathbb{Z}_{>0}$. We call e the *ramification index* of E/F .
- Noting that $k_F = \mathcal{O}_F/\mathfrak{p}_F$ is regarded as a subfield $k_E = \mathcal{O}_E/\mathfrak{p}_E$, we let f be the degree of the finite extension k_E/k_F . We call f the *residue degree* of E/F .

Note that these invariants satisfies the chain rule: if E/F is a finite separable extension with ramification index e and residue degree f and L/E is a finite separable extension with ramification index e' and residue degree f' , then L/F is a finite separable extension with ramification index ee' and residue degree ff' ,

Fact 2.8. *We have $n = ef$.*

- Definition 2.9.**
- (1) We say that E/F is *unramified* if $e = 1$ and (so, equivalently, $n = f$) and the residual extension k_E/k_F is separable.
 - (2) We say that E/F is *ramified* if it is not unramified.
 - (3) We say that E/F is *totally ramified* if $e = n$ (so, equivalently, $f = 1$).

Note we don't have to be worried about the second condition of the unramifiedness (separability of k_E/k_F) for local field since k_F is finite, hence perfect. Also, in this case, the ramifiedness is equivalent to that $e > 1$.

Example 2.10. Let p be an odd prime number such that $p \equiv -1 \pmod{4}$. Note that this condition is equivalent to that $\sqrt{-1} \notin \mathbb{F}_p$, which is furthermore equivalent to that $\sqrt{-1} \notin \mathbb{Q}_p$ by Hensel's lemma (explained later). We put $F_0 := \mathbb{Q}_p(\sqrt{-1})$ and $F_1 := \mathbb{Q}_p(\sqrt{p})$.

- The quadratic extension F_0/\mathbb{Q}_p is unramified since the residue field of F_0 must contain $\sqrt{-1}$, hence be a quadratic extension of \mathbb{F}_p .
- The quadratic extension F_1/\mathbb{Q}_p is ramified since the ring of integers \mathcal{O}_{F_1} contains \sqrt{p} and the ideal \mathfrak{p}_{E_1} generate by \sqrt{p} satisfies $\mathfrak{p}_{E_1}^2 = p\mathcal{O}_{F_1}$ (so \mathfrak{p}_{E_1} must be the maximal ideal).

In fact, unramified extensions are much easier to understand than ramified extensions. The fundamental reason for this lies in the following theorem:

Fact 2.11 (Hensel's lemma). *Let \mathcal{O} be a CDVR with maximal ideal \mathfrak{p} and residue field k . Let $f(X) \in \mathcal{O}[X]$ be a polynomial with mod \mathfrak{p} reduction $\bar{f}(X) \in k[X]$. If $\bar{\alpha} \in k$ is a simple root of $\bar{f}(X)$, then there uniquely exists a root $\alpha \in \mathcal{O}$ of $f(X)$ such that $\alpha \equiv \bar{\alpha} \pmod{\mathfrak{p}}$.*

Example 2.12. Let p be an odd prime number. Then \mathbb{Q}_p contains $\sqrt{-1}$ if and only if $p \equiv 1 \pmod{4}$. Indeed, note that the monic $X^2 + 1$ has a root in \mathbb{Q}_p if and only if it has a root in \mathbb{Z}_p since \mathbb{Z}_p is integrally closed. By Hensel's lemma, the latter condition is equivalent to that $X^2 + 1$ has a root in \mathbb{F}_p . Since $\sqrt{-1}$ is a primitive 4th root of unity (this is nothing but the definition of the symbol " $\sqrt{-1}$ ") and \mathbb{F}_p^\times is cyclic of order $p - 1$, we have $\sqrt{-1} \in \mathbb{F}_p^\times$ if and only if $4 \mid (p - 1)$, which means that $p \equiv 1 \pmod{4}$.

Proposition 2.13. *Let F be a CDVF with residue field k_F . The association $E \mapsto k_E$ for any finite unramified extension E/F gives a bijective map between the set of finite unramified extensions of F (in \bar{F}) and the set of finite separable extensions of k_F (in \bar{k}_F). Moreover, E/F is Galois if and only if so is k_E/k_F ; in this case the Galois groups are identified.*

Proof. We just give a sketch here. For checking the surjectivity, we take a finite separable extension k' of k_F . We write $k' = k_f[X]/(f(X))$ with $\bar{f}(X) \in k[X]$ and choose a lift $f(X) \in \mathcal{O}_F[X]$ of $\bar{f}(X)$. Then we can show that $F[X]/(f(X))$ is a finite unramified extension whose residue field is isomorphic to k' .

To show the remaining part, we take a finite unramified extension E of F . For the residual extension k_E/k_F , we choose $\bar{f}(X) \in k_F[X]$ as in the previous paragraph and lift it to $f(X) \in \mathcal{O}_F[X]$. Then, for any finite unramified extension E' , we have

$$\mathrm{Hom}_F(E, E') \xleftarrow{1:1} \mathrm{Hom}_{\mathcal{O}_F}(\mathcal{O}_E, \mathcal{O}_{E'}) \xleftarrow{1:1} \{\text{roots of } f(X) \text{ in } \mathcal{O}_{E'}\}$$

(if $\alpha' \in \mathcal{O}_{E'}$ is a root of $f(X)$, then the corresponding \mathcal{O}_F -algebra homomorphism is determined by $\alpha \mapsto \alpha'$). On the other hand, we also have

$$\mathrm{Hom}_{k_F}(k_E, k_{E'}) \xleftarrow{1:1} \{\text{roots of } \bar{f}(X) \text{ in } k_{E'}\}$$

By Hensel's lemma, the right-hand sides of these are naturally bijective. Thus we get a natural bijection $\mathrm{Hom}_F(E, E') \cong \mathrm{Hom}_{k_F}(k_E, k_{E'})$. This shows the injection of the map in the assertion. Also, being Galois is preserved between E/F and k_E/k_F . \square

Note that, in particular, when E and E' are finite unramified extensions of F , their composite field EE' is also a finite unramified extension of F ; this is the field corresponding to $k_E k_{E'}$ in the above proposition. Hence it makes sense to think about the *maximal unramified extension* of F , which is the direct limit (union) of all finite unramified extensions of F and denoted by F^{ur} . Then F^{ur} is a Galois extension of F whose Galois group $\mathrm{Gal}(F^{\mathrm{ur}}/F)$ is isomorphic to $\mathrm{Gal}(k_F^{\mathrm{sep}}/k_F)$. We remark that, for any finite extension E/F , the intersection $E \cap F^{\mathrm{ur}}$ gives the maximal unramified (over F) subextension of F in E ; in other words, $E/E \cap F^{\mathrm{ur}}$ is totally ramified and $E \cap F^{\mathrm{ur}}/F$ is unramified.

Let us apply this to the case of nonarchimedean local field. Let F be a nonarchimedean local field, hence k_F is a finite field, say \mathbb{F}_q (a field of q elements). As long as we fix an algebraic closure $\bar{\mathbb{F}}_q$ of \mathbb{F}_q , there uniquely exists a degree n extension of \mathbb{F}_q in $\bar{\mathbb{F}}_q$ for each $n \in \mathbb{Z}_{>0}$; it is \mathbb{F}_{q^n} , which is realized as the set of solutions of $x^{q^n} - x = 0$. This degree n extension $\mathbb{F}_{q^n}/\mathbb{F}_q$ is cyclic; $\mathrm{Gal}(\mathbb{F}_{q^n}/\mathbb{F}_q)$ has a natural generator called the *arithmetic Frobenius* element

$$\mathbb{F}_{q^n} \xrightarrow{\cong} \mathbb{F}_{q^n}; \quad x \mapsto x^q.$$

Note that the inverse to the arithmetic Frobenius element is also a generator. We call it the *geometric Frobenius* element and write $\mathrm{Frob}_{\mathbb{F}_q}$ for it⁴. Therefore, the Galois group of the infinite Galois extension $\bar{\mathbb{F}}_q/\mathbb{F}_q$ is isomorphic to the profinite completion $\hat{\mathbb{Z}}$ of \mathbb{Z} :

$$\mathrm{Gal}(\bar{\mathbb{F}}_q/\mathbb{F}_q) \cong \varprojlim_n \mathrm{Gal}(\mathbb{F}_{q^n}/\mathbb{F}_q) \cong \varprojlim_n \mathbb{Z}/n\mathbb{Z} =: \hat{\mathbb{Z}}.$$

Here, the topological generator 1 of $\hat{\mathbb{Z}}$ on the right-hand side corresponds to the arithmetic Frobenius element $\bar{\mathbb{F}}_q \xrightarrow{\cong} \bar{\mathbb{F}}_q: x \mapsto x^q$ on the left-hand side.

Now, by Proposition 2.13, for each $n \in \mathbb{Z}_{>0}$, there uniquely exists a degree n unramified extension F_n of F ; it is generated by the solutions to the equation $x^{q^n} - x = 0$. In other words, F_n is obtained by adjoining all $(q^n - 1)$ -th roots of unity to F .


Exercise 2.14. Let F be a nonarchimedean local field with residue field k_F of characteristic p . Prove that the maximal unramified extension F^{ur} is generated over F by roots of unity whose orders are prime-to- p .

We next consider ramified extensions. As mentioned before, ramified extensions are not so easy compared with unramified extension. For example, totally ramified extensions are not closed under the composition. Thus it does not make sense to think about something

⁴Here we have some conflict of notations: in Week 1, I used this symbol for denoting (a lift of) the arithmetic Frobenius.

like “maximal totally ramified extension”. Related to this, there is also no canonical way of associating a “maximal totally ramified subextension” to a given extension E/F .

Example 2.15. Let p , $F_0 = \mathbb{Q}_p(\sqrt{-1})$, and $F_1 := \mathbb{Q}_p(\sqrt{p})$ be as in Example 2.10. We furthermore introduce another quadratic extension $F_2 := \mathbb{Q}_p(\sqrt{-p})$, which is ramified for the same reason as F_1 . If we let E be the quartic extension $\mathbb{Q}_p(\sqrt{p}, \sqrt{-1})$ of \mathbb{Q}_p , then we have $E = F_0F_1 = F_0F_2 = F_1F_2$. The situation is summarized as follows:



In particular, note that the composite of two totally ramified extensions F_1 and F_2 contains an unramified extension.

Definition 2.16. Let \mathcal{O} be a CDVR with discrete valuation v . Let $f(X) \in \mathcal{O}[X]$ be a monic polynomial $f(X) = X^n + a_{n-1}X^{n-1} + \cdots + a_0$. We say that $f(X)$ is an *Eisenstein polynomial* if $v(a_i) \geq 1$ for any $1 \leq i \leq n-1$ and $v(a_0) = 1$.

Fact 2.17. Let \mathcal{O} be a CDVR with fractional field F . Let $f(X) \in \mathcal{O}[X]$ be an Eisenstein polynomial of degree n . Then $f(X)$ is irreducible and the field $F[X]/(f(X))$ is a totally ramified extension of F of degree n .

Exercise 2.18. Let $M_n(\mathbb{Q}_p)$ be the \mathbb{Q}_p -algebra of n -by- n matrices whose entries are in \mathbb{Q}_p . We consider the following element

$$\varphi := \begin{pmatrix} 0 & 1 & & & \\ & \ddots & \ddots & & \\ & & \ddots & \ddots & \\ & & & \ddots & 1 \\ p & & & & 0 \end{pmatrix} \in M_n(\mathbb{Q}_p).$$

More precisely, $(i, i+1)$ -entry of φ is 1 for $1 \leq i \leq n-1$, $(n, 1)$ -entry of φ is p , and all the other entries are 0. We consider the \mathbb{Q}_p -subalgebra $\mathbb{Q}_p[\varphi]$ of $M_n(\mathbb{Q}_p)$ generated by φ . Prove that $\mathbb{Q}_p[\varphi]$ is a finite extension of \mathbb{Q}_p (in particular, $\mathbb{Q}_p[\varphi]$ is a field). Also, determine the extension degree, the ramification index, and the residue degree of $\mathbb{Q}_p[\varphi]/\mathbb{Q}_p$.

2.3. Galois groups and Weil groups of local fields. Let E/F be a finite Galois extension of nonarchimedean local fields. Then, any element of $\text{Gal}(E/F)$ induces an element of the extension of residue fields k_E/k_F . In other words, we have a natural surjection $\text{Gal}(E/F) \twoheadrightarrow \text{Gal}(k_E/k_F)$. By letting E run over all finite Galois extensions of F , we also get a natural surjection $\Gamma_F := \text{Gal}(F^{\text{sep}}/F) \twoheadrightarrow \text{Gal}(\bar{k}_F/k_F)$.

Definition 2.19. We let I_F be the kernel of the map $\Gamma_F \twoheadrightarrow \text{Gal}(\bar{k}_F/k_F)$ and call it the *inertia subgroup* of Γ_F .

Recall that we have $\text{Gal}(F^{\text{ur}}/F) \cong \text{Gal}(\bar{k}_F/k_F)$. Hence the inertia subgroup I_F is nothing but the closed subgroup of Γ_F corresponding to the subextension F^{ur} , i.e., $I_F = \text{Gal}(F^{\text{sep}}/F^{\text{ur}})$.

Definition 2.20. We define a subgroup W_F of Γ_F to be the preimage of $\langle \text{Frob}_{k_F} \rangle$ under the map $\Gamma_F \twoheadrightarrow \text{Gal}(\bar{k}_F/k_F)$ and call it the *Weil group* of F .

$$\begin{array}{ccccccc} 1 & \longrightarrow & I_F & \longrightarrow & \Gamma_F & \longrightarrow & \text{Gal}(\bar{k}_F/k_F) \cong \hat{\mathbb{Z}} \longrightarrow 1 \\ & & \parallel & & \cup & & \cup \\ 1 & \longrightarrow & I_F & \longrightarrow & W_F & \longrightarrow & \langle \text{Frob}_{k_F} \rangle \cong \mathbb{Z} \longrightarrow 1 \end{array}$$

Note that the Weil group is not the Galois group for any Galois extension, hence there is no intrinsic topology on W_F . We equip W_F with the topology such that I_F is open in W_F and the induced topology on I_F coincides with the natural topology of I_F (as the Galois group of $F^{\text{sep}}/F^{\text{ur}}$). The natural inclusion $W_F \hookrightarrow \Gamma_F$ induces an inclusion between their maximal abelian quotients $W_F^{\text{ab}} \hookrightarrow \Gamma_F^{\text{ab}}$.

2.4. Local class field theory.

Theorem 2.21 (local class field theory). *Let F be a non-archimedean local field with residue field k . Then there uniquely exists an isomorphism*

$$\text{Art}_F: F^\times \xrightarrow{\cong} W_F^{\text{ab}}$$

as topological groups satisfying the following properties:

- (1) For any uniformizer $\varpi \in F^\times$, its image $\text{Art}_F(\varpi) \in W_F^{\text{ab}}$ is a lift of the geometric Frobenius $\text{Frob}_k \in \text{Gal}(\bar{k}/k)$.
- (2) For any finite separable extension E/F , the following diagram commutes:

$$\begin{array}{ccc} E^\times & \xrightarrow{\text{Art}_E} & W_E^{\text{ab}} \\ \text{Nr}_{E/F} \downarrow & & \downarrow \text{res} \\ F^\times & \xrightarrow{\text{Art}_F} & W_F^{\text{ab}} \end{array}$$

- (3) For any finite abelian extension E/F , Art_F induces an isomorphism

$$F^\times / \text{Nr}_{E/F}(E^\times) \xrightarrow{\cong} \text{Gal}(E/F).$$

Because of this theorem, it is important to know the structure of F^\times . So let us explain how F^\times can be understood.

We first note the exact sequence

$$1 \rightarrow \mathcal{O}_F^\times \rightarrow F^\times \xrightarrow{v} \mathbb{Z} \rightarrow 1.$$

This splits by choosing a uniformizer $\varpi \in F^\times$, i.e., we have $F^\times \cong \mathcal{O}_F^\times \times \langle \varpi \rangle$. Secondly, we have the exact sequence

$$1 \rightarrow (1 + \mathfrak{p}_F) \rightarrow \mathcal{O}_F^\times \rightarrow k_F^\times \rightarrow 1.$$

This splits by Hensel's lemma; elements of k_F^\times are identified with $(q-1)$ -roots of unity, where $q = |k_F|$. Finally, we consider the exponential/logarithm map between F and F^\times . Here, for simplicity, we suppose that $F = \mathbb{Q}_p$:

$$\begin{aligned} \exp: \mathbb{Q}_p^\times &\rightarrow \mathbb{Q}_p^\times; & x &\mapsto \sum_{n=0}^{\infty} \frac{1}{n!} x^n, \\ \log: \mathbb{Q}_p^\times &\rightarrow \mathbb{Q}_p; & x &\mapsto \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} (x-1)^n. \end{aligned}$$

These maps do not converge on the whole domain, but gives group isomorphisms between

$$\begin{cases} p\mathbb{Z}_p \text{ and } 1 + p\mathbb{Z}_p & \text{if } p \neq 2, \\ 4\mathbb{Z}_2 \text{ and } 1 + 4\mathbb{Z}_2 & \text{if } p = 2. \end{cases}$$

In the case where $p = 2$, we have $(1 + 2\mathbb{Z}_2) \cong \{\pm 1\} \times (1 + 4\mathbb{Z}_2)$. Thus, in conclusion, we have

$$\mathbb{Q}_p^\times \cong \begin{cases} \mathbb{Z} \times \mathbb{Z}/(p-1)\mathbb{Z} \times \mathbb{Z}_p & \text{if } p \neq 2, \\ \mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}_2 & \text{if } p = 2. \end{cases}$$

Exercise 2.22. Count the number of (isomorphism classes of) quadratic extensions of \mathbb{Q}_p .

Exercise 2.23. For any finite abelian group G , can we always find a finite abelian extension of nonarchimedean local fields E/F whose Galois group is isomorphic to G ?

3. WEEK 3: REPRESENTATIONS OF LOCALLY PROFINITE GROUPS

3.1. Locally profinite groups. The reference for this section is [BH06, §1].

- Definition 3.1.** (1) We say that a topological group G is *profinite* if G is compact and the set of open subgroups of G forms a fundamental system of neighborhood of $1 \in G$ (i.e., any open neighborhood of $1 \in G$ contains an open subgroup of G).
- (2) We say that a topological group G is *locally profinite* if it contains an open subgroup which is a profinite group.

Fact 3.2. A topological group G is profinite if and only if it is written as the inverse limit $G = \varprojlim_n G_n$ with respect to a projective system $\{G_n\}_n$ of finite groups.

We don't review here the fundamental properties of (locally) profinite groups, but just mention the following one, which will be used implicitly many many times.

Lemma 3.3. Let G be a profinite group. Then any open subgroup of H is of finite index.

Proof. Let us write $G = \bigsqcup_{g \in G/H} gH$. Then the g -translation $G \rightarrow G: x \mapsto gx$ is a homeomorphism, gH is also open in G . Thus, by the compactness of G , we conclude that the disjoint open covering $\{gH\}_{g \in G/H}$ must be a finite covering. Hence G/H is finite. \square

Example 3.4. (1) Any non-archimedean local field F is a locally profinite group as an additive group. Indeed, by the definition of its metric, the descending filtration

$$\mathcal{O}_F \supset \mathfrak{p}_F \supset \mathfrak{p}_F^2 \supset \cdots \supset \{0\}$$

consisting of open subgroups \mathfrak{p}^n forms a fundamental system of neighborhoods of 0. Since \mathcal{O}_F is closed (and also open) in F and bounded with respect to the metric, \mathcal{O}_F is compact (hence profinite). Note that we can write $\mathcal{O}_F \cong \varprojlim_n \mathcal{O}_F / \mathfrak{p}_F^n$.

(2) For any non-archimedean local field F , its multiplicative group F^\times is a locally profinite group. Indeed, by the definition of its metric, the descending filtration

$$\mathcal{O}_F^\times \supset (1 + \mathfrak{p}_F) \supset (1 + \mathfrak{p}_F^2) \supset \cdots \supset \{0\}$$

consisting of open subgroups $(1 + \mathfrak{p}^n)$ forms a fundamental system of neighborhoods of 1. Since \mathcal{O}_F^\times is closed (and also open) in F and bounded with respect to the metric, \mathcal{O}_F^\times is compact (hence profinite). Note that we can write $\mathcal{O}_F^\times \cong \varprojlim_n \mathcal{O}_F^\times / (1 + \mathfrak{p}_F^n)$.

(3) The previous examples can be generalized as follows.

For any non-archimedean local field F , the additive group $M_n(F)$ of n -by- n matrices is a locally profinite group. Here, we just regard $M_n(F)$ as $F^{\oplus n^2}$ and equipped it with the product topology. A fundamental system of its open neighborhood (of the zero matrix) can be taken to be

$$M_n(\mathcal{O}_F) \supset M_n(\mathfrak{p}_F) \supset M_n(\mathfrak{p}_F^2) \supset \cdots \supset \{0\}.$$

Next, we consider $G = \mathrm{GL}_n(F)$ for a non-archimedean local field F . Then, with respect to the induced topology from $M_n(F)$, G is a locally profinite group. A fundamental system of its open neighborhood (of the identity matrix) can be taken to be

$$\mathrm{GL}_n(\mathcal{O}_F) \supset 1 + M_n(\mathfrak{p}_F) \supset 1 + M_n(\mathfrak{p}_F^2) \supset \cdots \supset \{I_n\}.$$

(4) The previous example can be furthermore generalized as follows. Let \mathbf{G} be a linear algebraic group over F . Then, by definition, we can find an embedding (Zariski closed immersion) $\iota: \mathbf{G} \hookrightarrow \mathrm{GL}_n$ into some GL_n over F . Hence we may regard \mathbf{G} as a Zariski closed subgroup of GL_n via ι . Here, recall that ‘‘Zariski closed’’

means that \mathbf{G} can be defined to be the subset of GL_n consisting of zeros of some polynomials. Thus any element $g \in \mathrm{GL}_n(F) \setminus \mathbf{G}(F)$ is not a solution to those polynomials; then any element $h \in \mathrm{GL}_n(F) \setminus \mathbf{G}(F)$ which is “sufficiently” close to g in the locally profinite topology (p -adic topology) cannot be a solution. In other words, the complement $\mathrm{GL}_n(F) \setminus \mathbf{G}(F)$ is open, hence $\mathbf{G}(F)$ is closed in $\mathrm{GL}_n(F)$. In general, any closed subgroup of a locally profinite group is again a locally profinite group, hence so is $\mathbf{G}(F)$.

3.2. Smooth representations of locally profinite groups. The reference for this section is [BH06, §2].

Let G be a locally profinite group. In the following, by “a representation (π, V) ” of G , we mean a \mathbb{C} -vector space V equipped with an action π of G , i.e., a group homomorphism $\pi: G \rightarrow \mathrm{GL}_{\mathbb{C}}(V)$. (Sometimes we just say “ π is a representation of G ”.)

Definition 3.5. Let (π, V) be a representation of G . We say that (π, V) is *smooth* if any $v \in V$ is contained in V^K for some open compact subgroup K of G . Here, V^K denotes the subspace of K -fixed vectors, i.e.,

$$V^K := \{w \in V \mid \pi(k)(w) = w \text{ for any } k \in K\}.$$

In other words, (π, V) is smooth if and only if we have

$$V = \bigcup_{K \subset G} V^K,$$

where the index set is over all open compact subgroups K of G .

We want to examine examples of smooth representations. In representation theory of finite groups, an operation called *induction* plays a very important role in constructing representations of a given group. So let us consider whether the same technique is available in this context.

Let $H \subset G$ be a subgroup. What we want to do here is to construct a smooth representation of a “bigger” locally profinite group using a smooth representation of a “smaller” locally profinite group. So, firstly, let us assume that H is closed because this guarantees that H is again locally profinite. Let (σ, W) be a smooth representation of H . Let (π, V) be the induction of (σ, W) in the usual sense. More precisely, the underlying space V is

$$\{f: G \rightarrow W \mid f(hg) = \sigma(h)(f(g)) \text{ for any } h \in H\}$$

and the action π of G on V is given by the right-translation on the functions, i.e.,

$$(\pi(x)f)(g) := f(gx).$$

Then, is (π, V) smooth? In fact, NO in general. So that (π, V) is smooth, for any f , there must be an open compact subgroup $K \subset G$ satisfying $f(gK) = f(g)$ (for any $g \in G$). However, this property is not formally deduced from the definition of V in general.

The idea is to modify the definition of V so that this condition is satisfied. In other words, if we put

$$V^\infty := \{f: G \rightarrow W \mid \exists K \text{ s.t. } f(hgk) = \sigma(h)(f(g)) \text{ for any } h \in H, k \in K\},$$

then (π, V^∞) is smooth (with respect to the same right-translation action π).

Definition 3.6. Let H be a closed subgroup of G and (σ, W) be a smooth representation of H . We call the smooth representation (π, V) defined as in the previous paragraph the *smooth induction* of (σ, W) from H to G . In our context, we always only consider the smooth induction, so we just say the *induction* of (σ, W) and write $(\mathrm{Ind}_H^G \sigma, \mathrm{Ind}_H^G W)$ for it.

Before we think about examples, let us introduce one guiding fact:

Fact 3.7. *Any irreducible smooth representation of $\mathrm{GL}_n(F)$ is either one-dimensional (i.e., a character) or infinite dimensional.*

This fact says that, in representation theory of locally profinite groups, we seriously have to face infinite dimensional representations. However, it is still possible to formulate some finiteness condition for smooth representations; it is called the *admissibility*.

Definition 3.8. Let (π, V) be a representation of G . We say that (π, V) is *admissible* if it is smooth and $\dim_{\mathbb{C}}(V^K)$ is finite-dimensional for any open compact subgroup K of G .

Example 3.9. Let $G = \mathrm{GL}_2(F)$, where F is a nonarchimedean local field.

- (1) Let $\chi: G \rightarrow \mathbb{C}^\times$ be a character, or equivalently, one-dimensional representation (χ, \mathbb{C}) . Then, by definition, (χ, \mathbb{C}) is smooth if and only if χ is trivial on some open compact subgroup of G . (This is equivalent to that χ is continuous with respect to the discrete topology of \mathbb{C}^\times .) Any smooth character of G is of course admissible.
- (2) Let $B \subset G$ be the subgroup of upper-triangular matrices (this is a closed subgroup). Let (π, V) be the (smooth) induction $(\mathrm{Ind}_B^G \mathbb{1}, \mathrm{Ind}_B^G \mathbb{C})$ of the trivial representation $(\mathbb{1}, \mathbb{C})$ of B to G . By definition, we can explicitly write

$$V = \{f: B \backslash G \rightarrow \mathbb{C} \mid \exists K \text{ s.t. } f(gk) = f(g) \text{ for any } k \in K\}.$$

To check the admissibility of (π, V) , let us fix any open compact subgroup K of G . Then we have

$$V^K \cong \{f: B \backslash G/K \rightarrow \mathbb{C}\}.$$

This is finite-dimensional since $B \backslash G/K$ is finite. Indeed, to check it, we may replace K with any its open subgroup freely (recall that such a subgroup must be of finite index in K). Especially, we may assume that K is contained in $\mathrm{GL}_2(\mathcal{O}_F)$. Since K must be also open, hence of finite index, in $\mathrm{GL}_2(\mathcal{O}_F)$, it is enough to show that $B \backslash G/\mathrm{GL}_2(\mathcal{O}_F)$ is a finite set. It is a well-known fact that $G = B \mathrm{GL}_2(\mathcal{O}_F)$ (so-called the *Iwasawa decomposition*), hence $B \backslash G/\mathrm{GL}_2(\mathcal{O}_F)$ is a singleton.

- (3) Next consider the subgroup $T \subset G$ of diagonal matrices (this is a closed subgroup). Let (π, V) be the (smooth) induction $(\mathrm{Ind}_T^G \mathbb{1}, \mathrm{Ind}_T^G \mathbb{C})$ of the trivial representation $(\mathbb{1}, \mathbb{C})$ of T to G . By definition, we can explicitly write

$$V = \{f: T \backslash G \rightarrow \mathbb{C} \mid \exists K \text{ s.t. } f(gk) = f(g) \text{ for any } k \in K\}.$$

To check the admissibility of (π, V) , let us fix any open compact subgroup K of G . Then we have

$$V^K \cong \{f: T \backslash G/K \rightarrow \mathbb{C}\}.$$

However, this space is infinite dimensional (Exercise below). Hence (π, V) is smooth but not admissible.

Note that this example shows that B is large enough so that the admissibility is preserved by the induction to G , but T is too small. This idea will be elaborated as the “parabolic induction” later.

Exercise 3.10. Prove that the set $T \backslash G/K$ in the above example is infinite.

Fact 3.11. *Let $G = \mathbf{G}(F)$ for any connected reductive group \mathbf{G} over a nonarchimedean local field F . Then any irreducible smooth representation of G is admissible.*

3.3. Frobenius reciprocity. Recall that, in representation theory of finite groups, we have so-called the *Frobenius reciprocity*, which is the adjunction between the induction functor and the restriction functor. In fact, the Frobenius reciprocity holds also for the smooth induction as well.

Theorem 3.12 (Frobenius reciprocity for Ind). *Let H be a closed subgroup of G and (σ, W) a smooth representation of G . Let (π, V) be a smooth representation of G . Then we have an isomorphism*

$$\mathrm{Hom}_G(\pi, \mathrm{Ind}_H^G \sigma) \cong \mathrm{Hom}_H(\pi|_H, \sigma).$$

given by $\phi \mapsto \alpha \circ \phi$, where $\alpha: \mathrm{Ind}_H^G \sigma \rightarrow \sigma$ is $f \mapsto f(1)$.

For the proof, see [BH06, §2.4].

Here, we caution that the smooth induction is put on the target in $\mathrm{Hom}(-, -)$. In other words, the smooth induction is the right adjoint to the restriction. In contrast to the case of finite groups, representations may not be semisimple. Thus we cannot swap the source and target in $\mathrm{Hom}(-, -)$ freely in general.

Then, does the restriction have a left adjoint? In fact, the answer is YES when H is open; it is given by the following variant of a smooth induction:

Definition 3.13. Let H be an open subgroup of G and (σ, W) be a smooth representation of H . We put

$$V_c^\infty := \left\{ f: G \rightarrow W \left| \begin{array}{l} \bullet f \text{ is compactly supported modulo } H \\ \bullet \exists K \text{ s.t. } f(hgk) = \sigma(h)(f(g)) \text{ for any } h \in H, k \in K \end{array} \right. \right\}$$

and consider the right-translation action π of G on V_c^∞ . Then (π, V_c^∞) is a smooth representation of G . We call it the *compact induction* of (σ, W) from H to G and write $(\mathrm{c}\text{-Ind}_H^G \sigma, \mathrm{c}\text{-Ind}_H^G W)$.

Theorem 3.14 (Frobenius reciprocity for c-Ind). *Let H be a closed subgroup of G and (σ, W) a smooth representation of G . Let (π, V) be a smooth representation of G . Then we have an isomorphism*

$$\mathrm{Hom}_G(\mathrm{c}\text{-Ind}_H^G \sigma, \pi) \cong \mathrm{Hom}_H(\sigma, \pi|_H).$$

given by $\phi \mapsto \phi \circ \beta$. Here, $\beta: \sigma \mapsto \mathrm{c}\text{-Ind}_H^G \sigma$ is $w \mapsto f_w$, where $f_w: H \rightarrow W$ is such that $f_w(h) = \sigma(h)w$ for $h \in H$ and $f_w(g) = 0$ for $g \in G \setminus H$.

3.4. Representations of profinite groups. We define the notion of a subrepresentation and a direct sum of smooth representations in the usual way. The following proposition is a simple consequence of Zorn's lemma (see [BH06, §2.2, Proposition]).

Proposition 3.15. *For any smooth representation (π, V) , the following are equivalent:*

- (1) (π, V) is the direct sum of irreducible subrepresentations.
- (2) (π, V) is the sum of irreducible subrepresentations.
- (3) for any subrepresentation (π_1, V_1) , there exists a complement, i.e., another subrepresentation (π_2, V_2) such that $(\pi, V) \cong (\pi_1, V_1) \oplus (\pi_2, V_2)$.

Definition 3.16. When a smooth representation satisfies the conditions in the previous proposition, we say it is *semisimple*.

Note that, in contrast to the case of finite groups, there exist plenty non-semisimple smooth representations in general.

Exercise 3.17. Show that $\text{Ind}_B^G \mathbb{1}$ in Example 3.9 (2) is non-semisimple. (Hint: check that $\text{Ind}_B^G \mathbb{1}$ is not irreducible by finding a proper subrepresentation and show that $\text{End}_G(\text{Ind}_B^G \mathbb{1}) = 1$ using Frobenius reciprocity and Schu'r lemma, which will be explained later.)

Proposition 3.18. *Any smooth representation of a profinite group K is semisimple.*

Proof. Let (π, V) be a smooth representation of K . Then, for each $v \in V$, there exists an open (hence of finite index) subgroup K' of K such that $v \in V^{K'}$. Here, by shrinking K' if necessary, we may assume that K' is normal in K . (Indeed, by writing $K = \bigcup_{l \in K/K'} lK'$, it is enough to replace K' with $\bigcap_{l \in K/K'} lK'l^{-1}$.) Note that v is contained in the subrepresentation $\text{Span}_{\mathbb{C}}\{\pi(k)v \mid k \in K\}$ generated by v . However, since the action of K' on this subrepresentation is trivial and K/K' is a finite group, this subrepresentation must be semisimple. Thus, (π, V) can be written as a sum of semisimple representations, hence so is itself. \square

Definition 3.19. Let (π, V) be a smooth representation of G . For any open compact subgroup K of G , by the previous proposition, we can write

$$V = \bigoplus_{\sigma \in \text{Irr}(K)} V[\sigma].$$

Here, the both hand sides are regarded as representations of K and $V[\sigma]$ denotes the sum of irreducible K -subrepresentations of V isomorphic to σ ($\text{Irr}(K)$ is the set of isomorphism classes of irreducible smooth representations of K). We call $V[\sigma]$ the σ -isotypic part of V . Note that $V^K = V[\mathbb{1}]$.

3.5. Contragredient representation. Recall that, for any representation (π, V) of a finite group G , its dual (*contragredient*) representation (π^*, V^*) is defined by $V^* := \text{Hom}_{\mathbb{C}}(V, \mathbb{C})$ and

$$\langle \pi^*(g)(v^*), v \rangle = \langle v^*, \pi(g^{-1})(v) \rangle$$

for any $g, v \in V, v^* \in V^*$. In the context of locally profinite groups, this definition contains the issue as in the definition of Ind ; i.e., the resulting representation may not be smooth. So, again, we consider smoothening V^* .

Definition 3.20. For a smooth representation (π, V) of G , we define its *contragredient* representation (π^\vee, V^\vee) by

$$V^\vee := \bigcup_{K \subset G} (V^*)^K,$$

where K runs open compact subgroups of G and $\pi^\vee = \pi^*|_{V^\vee}$.

Exercise 3.21. Show that, for any open compact subgroup K of G , we have $(V^\vee)^K \cong (V^K)^*$.

Proposition 3.22. *For a smooth representation (π, V) of G , we consider the natural map $\pi \rightarrow (\pi^\vee)^\vee$ given by $v \mapsto [v^\vee \mapsto \langle v^\vee, v \rangle]$. This map is G -equivariant. Moreover, it is isomorphic if and only if (π, V) is admissible.*

Proof. The first statement can be checked easily. Then, for any open compact subgroup K of G , we get $\pi^K \rightarrow ((\pi^\vee)^\vee)^K$. Since $\pi = \bigcup_K \pi^K$ and $(\pi^\vee)^\vee = \bigcup_K ((\pi^\vee)^\vee)^K$, it is enough to discuss when this map is bijective (for all K). By applying the previous exercise twice, we see that this map is identified with the natural map $\pi^K \rightarrow ((\pi^K)^*)^*$. It is well-known fact in linear algebra that this natural map is bijective if and only if π^K is finite-dimensional. \square

3.6. Irreducible representations and Schur's lemma.

Definition 3.23. Let (π, V) be a smooth representation of G . We say that (π, V) is *irreducible* if (π, V) has no G -subspace (subrepresentation) other than $\{0\}$ and V .

Lemma 3.24. Let (π, V) be an irreducible smooth representation of G . Then V is generated by any nonzero vector $v \in V$, i.e., we have $V = \text{Span}_{\mathbb{C}}(\{\pi(g)v \mid g \in G\})$.

Proof. The \mathbb{C} -vector subspace $\text{Span}_{\mathbb{C}}(\{\pi(g)v \mid g \in G\})$ of V is stable under G -action. Thus the irreducibility of V implies that it equals $\{0\}$ or V . Since $v \neq 0$, it must be V . \square

Definition 3.25. For smooth representations (π, V) and (π', V') , we define the set $\text{Hom}_G(\pi, \pi')$ of G -equivariant homomorphisms by

$$\text{Hom}_G(\pi, \pi') := \{\phi \in \text{Hom}_{\mathbb{C}}(V, V') \mid \phi(\pi(g)v) = \pi'(g)\phi(v) \forall g \in G, \forall v \in V\}.$$

When $(\pi, V) = (\pi, V')$, we simply write $\text{End}_G(\pi)$ for $\text{Hom}_G(\pi, \pi)$.

Theorem 3.26 (Schur's lemma). Let (π, V) be an irreducible smooth representation of G . Assume that $\dim_{\mathbb{C}}(V)$ is countable. Then we have $\text{End}_G(\pi) = \mathbb{C}$.

Proof. Suppose that $\phi \in \text{End}_G(\pi)$ is a nonzero G -equivariant endomorphism of (π, V) . Then both $\text{Ker}(\phi)$ and $\text{Im}(\phi)$ are G -stable subspaces of V . Hence, by the irreducibility of V , they must be $\{0\}$ or V . Since ϕ is supposed to be nonzero, we necessarily have $\text{Ker}(\phi) = 0$ and $\text{Im}(\phi) = V$; in other words, ϕ is an isomorphism. Therefore, $\text{End}_G(\pi)$ is a division \mathbb{C} -algebra (i.e., possibly non-commutative \mathbb{C} -algebra whose any nonzero element is invertible).

By Lemma 3.24, if we fix any nonzero vector $v \in V$, then v generates V . Hence, any G -equivariant endomorphism $\phi \in \text{End}_G(\pi)$ is determined uniquely by the image $\phi(v)$ of v . If $\phi(v) \in V$ is equal to $\phi'(v) \in V$ up to scalar, then $\phi \text{End}_G(\pi)$ equals $\phi' \text{End}_G(\pi)$ up to scalar. In particular, the dimension of $\text{End}_G(\pi)$ as a \mathbb{C} -vector space is bounded by the dimension of V . Since $\dim_{\mathbb{C}}(V)$ is countable, so is $\dim_{\mathbb{C}}(\text{End}_G(\pi))$.

Now suppose that $\dim_{\mathbb{C}}(\text{End}_G(\pi))$ is bigger than \mathbb{C} ; then we can choose $\phi \in \text{End}_G(\pi) \setminus \mathbb{C}$. Then the division \mathbb{C} -algebra $\text{End}_G(\pi)$ contains the rational function field $\mathbb{C}(\phi)$. However, the dimension of $\mathbb{C}(\phi)$ as a \mathbb{C} -vector space is uncountable. (For example, it can be easily checked that the subset $\{(\phi - a)^{-1} \mid a \in \mathbb{C}\}$ is linear independent.) Thus we get a contradiction. \square

A reasonable sufficient condition for that the assumption of Theorem 3.26 is satisfied is the following:

Lemma 3.27. If there exists an open compact subgroup K_0 of G such that G/K_0 is countable, then any irreducible representation of G has countable dimension.

Proof. Note that if K_0 is an open compact subgroup whose G/K_0 is countable, then any open compact subgroup K satisfies that G/K is countable. (Indeed, since $K \cap K_0$ is also open subgroup of K_0 , it is compact and of finite index in K_0 . Thus $G/(K \cap K_0)$ is countable. As $K \cap K_0$ is also of finite index in K , G/K is countable.) By Lemma 3.24, any nonzero vector $v \in V$ generates V . If we let K be an open compact subgroup of G satisfying $v \in V^K$, then $\dim_{\mathbb{C}}(V)$ is bounded by the cardinality of G/K , which is countable. \square

Example 3.28. When $G = \text{GL}_n(F)$, G satisfies the countability assumption in the above lemma. Indeed, if we put $K_0 := \text{GL}_n(\mathcal{O}_F)$, then K_0 is an open compact subgroup of G . Moreover, we have the following decomposition (so-called “Cartan decomposition”, which is

a consequence of elementary divisor theory):

$$G = \bigsqcup_{\substack{a,b \in \mathbb{Z} \\ a \leq b}} K_0 \begin{pmatrix} \varpi^a & 0 \\ 0 & \varpi^b \end{pmatrix} K_0.$$

Each summand is compact, hence its right- K_0 -cosets are finite since K_0 is open. As the index set is countable, we only have countable many right- K_0 -cosets in G .

More generally, for any linear algebraic group \mathbf{G} over F , $G := \mathbf{G}(F)$ satisfies the countability assumption. (take an embedding $\mathbf{G} \hookrightarrow \mathrm{GL}_n$ and put $K_0 := G \cap \mathrm{GL}_n(\mathcal{O}_F)$).

In the following, we always assume that there exists an open compact subgroup K_0 whose G/K_0 is countable. Let us suppose that (π, V) is an irreducible representation of G . Let Z be the center of G . Then, for any $z \in Z$, the automorphism $\pi(z) \in \mathrm{GL}_{\mathbb{C}}(V)$ is G -equivariant. Indeed, for any $g \in G$ and $v \in V$, we have

$$\pi(z)(\pi(g)v) = \pi(zg)v = \pi(gz)v = \pi(g)(\pi(z)v).$$

By Schur's lemma, $\pi(z)$ must be a (nonzero) scalar multiplication. Therefore, we get a map $Z \rightarrow \mathbb{C}^\times$. It is easy to check that this map is a smooth character.

Definition 3.29. For any irreducible representation (π, V) of G , we call the smooth character of Z defined in this way *the central character of (π, V)* and write ω_π .

4. WEEK 4: IRREDUCIBLE SMOOTH REPRESENTATIONS OF $\mathrm{GL}_2(F)$

Let F be a non-archimedean local field of residual characteristic p , hence a finite extension of \mathbb{Q}_p or $\mathbb{F}_p((t))$. The aim of this week is to give an overview on a classification of irreducible smooth representations of group $\mathrm{GL}_2(F)$.

4.1. Recap on irreducible representations of $\mathrm{GL}_2(\mathbb{F}_q)$. Let \mathbb{F}_q be a finite field of order q and characteristic p . We first review how the irreducible representations of $\mathrm{GL}_2(\mathbb{F}_q)$ are classified.

Let us simply write $G = \mathrm{GL}_2$ in the following. We introduce several subgroups of $\mathrm{GL}_2(\mathbb{F}_q)$ as follows:

$$\begin{aligned} B(\mathbb{F}_q) &:= \left\{ \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \in M_2(\mathbb{F}_q) \mid a, d \in \mathbb{F}_q^\times, b \in \mathbb{F}_q \right\}, \\ T(\mathbb{F}_q) &:= \left\{ \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \in M_2(\mathbb{F}_q) \mid a, d \in \mathbb{F}_q^\times \right\}, \\ U(\mathbb{F}_q) &:= \left\{ \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \in M_2(\mathbb{F}_q) \mid b \in \mathbb{F}_q \right\}. \end{aligned}$$

Note that $U(\mathbb{F}_q)$ is a normal subgroup in $B(\mathbb{F}_q)$ and that we have the semi-direct decomposition $B(\mathbb{F}_q) = T(\mathbb{F}_q) \ltimes U(\mathbb{F}_q)$. In particular, we have a natural surjection $B(\mathbb{F}_q) \twoheadrightarrow T(\mathbb{F}_q)$ by quotienting by $U(\mathbb{F}_q) \triangleleft B(\mathbb{F}_q)$.

Let us take two characters χ_1, χ_2 of \mathbb{F}_q^\times . Then we get a character of $T(\mathbb{F}_q)$

$$\chi := \chi_1 \boxtimes \chi_2: T(\mathbb{F}_q) \rightarrow \mathbb{C}^\times; \quad \mathrm{diag}(t_1, t_2) \mapsto \chi_1(t_1) \cdot \chi_2(t_2).$$

By pulling back χ to $B(\mathbb{F}_q)$, we may regard χ as a character of $B(\mathbb{F}_q)$. Finally, by taking the induction to $G(\mathbb{F}_q)$, we get a representation $\mathrm{Ind}_{B(\mathbb{F}_q)}^{G(\mathbb{F}_q)} \chi$ of $G(\mathbb{F}_q)$. We call the representation $\mathrm{Ind}_{B(\mathbb{F}_q)}^{G(\mathbb{F}_q)} \chi$ a *principal series representation* (associated to χ).

The decomposition rule of $\mathrm{Ind}_{B(\mathbb{F}_q)}^{G(\mathbb{F}_q)} \chi$ is described as follows.

Proposition 4.1 ([BH06, Section 6]). *(1) When $\chi_1 \neq \chi_2$, $\mathrm{Ind}_{B(\mathbb{F}_q)}^{G(\mathbb{F}_q)} \chi$ is irreducible.*

(2) When $\chi_1 = \chi_2$ (say χ_0), we have $\mathrm{Ind}_{B(\mathbb{F}_q)}^{G(\mathbb{F}_q)} \chi \cong (\mathrm{Ind}_{B(\mathbb{F}_q)}^{G(\mathbb{F}_q)} \mathbb{1}) \otimes (\chi_0 \circ \det)$ and $\mathrm{Ind}_{B(\mathbb{F}_q)}^{G(\mathbb{F}_q)} \mathbb{1}$ decomposes into the sum of two irreducible representations; the one is the trivial representation $\mathbb{1}$ of $G(\mathbb{F}_q)$ and the other one is called the Steinberg representation St . In summary, we have $\mathrm{Ind}_{B(\mathbb{F}_q)}^{G(\mathbb{F}_q)} \chi = (\chi_0 \circ \det) \oplus \mathrm{St} \otimes (\chi_0 \circ \det)$.

(3) Two principal series representations $\mathrm{Ind}_{B(\mathbb{F}_q)}^{G(\mathbb{F}_q)} \chi_1 \boxtimes \chi_2$ and $\mathrm{Ind}_{B(\mathbb{F}_q)}^{G(\mathbb{F}_q)} \chi'_1 \boxtimes \chi'_2$ contains a common irreducible representation if and only if $(\chi_1, \chi_2) = (\chi'_1, \chi'_2), (\chi'_2, \chi'_1)$.

Can this construction exhaust all irreducible representations of $\mathrm{GL}_2(\mathbb{F}_q)$? In fact, not; the missing representations are called *cuspidal* representations. In my course of the previous semester [Oi24], I introduced two ways of constructing all cuspidal representations of $\mathrm{GL}_2(\mathbb{F}_q)$. The one is a purely algebraic construction; we first define a virtual representation as a linear combination of several induced representations, and then show that it is in fact an irreducible representation which is not contained in any principal series representation. The details of this construction can be found in [BH06, Section 6]. The other one is a geometric construction; we first define an algebraic variety on which $\mathrm{GL}_2(\mathbb{F}_q)$ acts and then take its ℓ -adic étale cohomology. Then the resulting cohomology realizes cuspidal representations (and even also principal series representations); this is what is called *Deligne–Lusztig theory* [DL76].

4.2. Principal series representations of $\mathrm{GL}_2(F)$. Now let us consider the group $\mathrm{GL}_2(F)$. Recall that, in the last week, we have introduced the notion of the (smooth) induction in the context of smooth representation theory of locally profinite groups. Thus it is natural to try to imitate the construction of principal series also for $\mathrm{GL}_2(F)$.

We introduce several subgroups of $\mathrm{GL}_2(F)$ in the same way as before, just by replacing \mathbb{F}_q with F :

$$B(F) := \left\{ \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \in M_2(F) \mid a, d \in F^\times, b \in F \right\},$$

$$T(F) := \left\{ \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \in M_2(F) \mid a, d \in F^\times \right\},$$

$$U(F) := \left\{ \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \in M_2(F) \mid b \in F \right\}.$$

Then these groups satisfy the same properties as before. Especially, $U(F)$ is a normal subgroup in $B(F)$ and we have the semi-direct decomposition $B(F) = T(F) \rtimes U(F)$. We have a natural surjection $B(F) \twoheadrightarrow T(F)$ by quotienting by $U(F) \triangleleft B(F)$.

Let us take two smooth characters χ_1, χ_2 of F^\times . Then we can define the representation $\mathrm{Ind}_{B(F)}^{G(F)} \chi_1 \boxtimes \chi_2$ of $\mathrm{GL}_2(F)$ in exactly the same manner as before. We call $\mathrm{Ind}_{B(F)}^{G(F)} \chi_1 \boxtimes \chi_2$ a *principal series representation* (associated to $\chi_1 \boxtimes \chi_2$). However, the decomposition rule of $\mathrm{Ind}_{B(F)}^{G(F)} \chi_1 \boxtimes \chi_2$ is a bit more difficult than the case of $\mathrm{GL}_2(\mathbb{F}_q)$. The point is that representations of $\mathrm{GL}_2(\mathbb{F}_q)$ are always semisimple, but those of $\mathrm{GL}_2(F)$ are not. To be more precise, the situation is summarized as follows.

Let \mathbb{F}_q be the residue field of F . Let $|\cdot|: F^\times \rightarrow \mathbb{C}^\times$ denote the absolute value character, i.e., $|x| = q^{-v(x)}$, where v is the normalized valuation of F .

Proposition 4.2 ([BH06, Section 9]). *(1) The representation $\mathrm{Ind}_{B(F)}^{G(F)} \chi_1 \boxtimes \chi_2$ is reducible if and only if $\chi_1 \chi_2^{-1}$ equals either $\mathbb{1}$ or $|\cdot|^2$. Moreover, $\mathrm{Ind}_{B(F)}^{G(F)} \chi_1 \boxtimes \chi_2$ and $\mathrm{Ind}_{B(F)}^{G(F)} \chi'_1 \boxtimes \chi'_2$ are isomorphic if and only if (χ_1, χ_2) equals (χ'_1, χ'_2) or $(\chi'_2 \cdot |\cdot|^{-\frac{1}{2}}, \chi'_1 \cdot |\cdot|^{\frac{1}{2}})$.*
(2) Suppose that $\chi_1 \chi_2^{-1} = \mathbb{1}$, hence $\chi_1 = \chi_2 = \chi_0$ for some smooth character $\chi_0: F^\times \rightarrow \mathbb{C}^\times$. Then $\mathrm{Ind}_{B(F)}^{G(F)} \chi_1 \boxtimes \chi_2 \cong \mathrm{Ind}_{B(F)}^{G(F)} \mathbb{1} \otimes (\chi_0 \circ \det)$ and we have the following non-split exact sequence of smooth representations of $\mathrm{GL}_2(F)$

$$0 \rightarrow \mathbb{1} \rightarrow \mathrm{Ind}_{B(F)}^{G(F)} \mathbb{1} \rightarrow \mathrm{St} \rightarrow 0,$$

where $\mathbb{1}$ is the trivial representation of $\mathrm{GL}_2(F)$ and St is an infinite-dimensional irreducible smooth representation of $\mathrm{GL}_2(F)$ (called the Steinberg representation). In other words, $\mathbb{1}$ is the unique irreducible subrepresentation of $\mathrm{Ind}_{B(F)}^{G(F)} \mathbb{1}$ which cannot be a quotient, and St is the unique irreducible quotient representation of $\mathrm{Ind}_{B(F)}^{G(F)} \mathbb{1}$ which cannot be a subrepresentation.

(3) Suppose that $\chi_1 \chi_2^{-1} = |\cdot|^2$, hence $\chi_1 = \chi_0 \cdot |\cdot|$ and $\chi_2 = \chi_0 \cdot |\cdot|^{-1}$ for some smooth character $\chi_0: F^\times \rightarrow \mathbb{C}^\times$. Then $\mathrm{Ind}_{B(F)}^{G(F)} \chi_1 \boxtimes \chi_2 \cong \mathrm{Ind}_{B(F)}^{G(F)} (|\cdot| \boxtimes |\cdot|^{-1}) \otimes (\chi_0 \circ \det)$ and we have the following non-split exact sequence of smooth representations of $\mathrm{GL}_2(F)$

$$0 \rightarrow \mathrm{St} \rightarrow \mathrm{Ind}_{B(F)}^{G(F)} (|\cdot| \boxtimes |\cdot|^{-1}) \rightarrow \mathbb{1} \rightarrow 0.$$

In other words, $\mathbb{1}$ is the unique irreducible quotient representation which cannot be a subrepresentation, and St is the unique irreducible subrepresentation which cannot be a quotient representation.

As in the case of $\text{GL}_2(\mathbb{F}_q)$, this construction does not produce all irreducible smooth representations of $\text{GL}_2(F)$. If an irreducible smooth representation of $\text{GL}_2(F)$ is not contained in any principal series representation, we call it a *supercuspidal representation*.

4.3. Depth-zero supercuspidal representations of $\text{GL}_2(F)$. The question is: how to construct (all) irreducible supercuspidal representations of $\text{GL}_2(F)$? For principal series representations, we just imitated the construction in the case of $\text{GL}_2(\mathbb{F}_q)$. However, for supercuspidal representations, we can immediately see that the construction in the finite field case no longer works.

One idea is, instead of imitating, “reducing” the construction to the finite field case. The point is that $\text{GL}_2(F)$ has the following open compact subgroup:

$$\text{GL}_2(\mathcal{O}) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(F) \mid a, b, c, d \in \mathcal{O}, ad - bc \in \mathcal{O}^\times \right\},$$

where \mathcal{O} denotes the ring of integers in F . When each entry of a 2-by-2 matrix belongs to \mathcal{O} , it makes sense to take its mod- \mathfrak{p} reduction for the maximal ideal \mathfrak{p} of \mathcal{O} . Then we get a 2-by-2 matrix with entries in $\mathbb{F}_q = \mathcal{O}/\mathfrak{p}$:

$$\text{GL}_2(\mathcal{O}) \rightarrow \text{GL}_2(\mathbb{F}_q): \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \begin{pmatrix} \bar{a} & \bar{b} \\ \bar{c} & \bar{d} \end{pmatrix}.$$

Exercise 4.3. Prove that the map $\text{SL}_2(\mathcal{O}) \rightarrow \text{SL}_2(\mathbb{F}_q)$ is also surjective.

Now let κ be an irreducible cuspidal representation of $\text{GL}_2(\mathbb{F}_q)$. By pulling back it along the above surjection, we get an irreducible smooth representation of $\text{GL}_2(\mathcal{O})$ (let’s again write κ). Recall that, in the last week, we introduced a variant of the usual (smooth) induction, which is called the compact induction “c-Ind”. The basic strategy is to construct a smooth representation of $\text{GL}_2(F)$ by applying the compact induction to this representation κ of $\text{GL}_2(\mathcal{O})$.

However, here we have an obvious obstruction. Let $Z(F)$ be the center of $\text{GL}_2(F)$, i.e., the subgroup of non-zero scalar matrices. Then we have

$$\text{c-Ind}_{\text{GL}_2(\mathcal{O})}^{\text{GL}_2(F)} \kappa = \text{c-Ind}_{Z(F)\text{GL}_2(\mathcal{O})}^{\text{GL}_2(F)} \left(\text{c-Ind}_{\text{GL}_2(\mathcal{O})}^{Z(F)\text{GL}_2(\mathcal{O})} \kappa \right).$$

Since the quotient $Z(F)\text{GL}_2(\mathcal{O})/\text{GL}_2(\mathcal{O})$ is isomorphic to $Z(F)/Z(\mathcal{O}) \cong F^\times/\mathcal{O}^\times$, which is an infinite abelian group, the internal induction $\text{c-Ind}_{\text{GL}_2(\mathcal{O})}^{Z(F)\text{GL}_2(\mathcal{O})} \kappa$ breaks into infinitely many pieces of irreducible representations of $Z(F)\text{GL}_2(\mathcal{O})$. So the whole compact induction cannot be irreducible.

To remedy this issue, we first extend the representation κ from $\text{GL}_2(\mathcal{O})$ to $Z(F)\text{GL}_2(\mathcal{O})$ by fixing a character of $Z(F)$. Note that, as κ is irreducible, the restriction of κ to $Z(\mathcal{O})$ is given by a character (“central character”). Let ω be a character of $Z(F)$ such that $\omega|_{Z(\mathcal{O})}$ coincides with the central character of κ . We define a representation $\tilde{\kappa}$ of $Z(F)\text{GL}_2(\mathcal{O})$ by

$$\tilde{\kappa}(z) := \begin{cases} \omega(z) & \text{if } z \in Z(F), \\ \kappa(g) & \text{if } g \in \text{GL}_2(\mathcal{O}). \end{cases}$$

We put

$$\pi_{\tilde{\kappa}} := \text{c-Ind}_{Z(F)\text{GL}_2(\mathcal{O})}^{\text{GL}_2(F)} \tilde{\kappa}.$$

Fact 4.4. *The representation $\pi_{\bar{\kappa}}$ is an irreducible supercuspidal representation of $\mathrm{GL}_2(F)$. Moreover, for any other κ' and ω' , the representations $\pi_{\bar{\kappa}}$ and $\pi_{\bar{\kappa}'}$ are isomorphic if and only if $\kappa \cong \kappa'$ and $\omega = \omega'$.*

Now we come up with the next question: does this construction exhaust all irreducible supercuspidal representations of $\mathrm{GL}_2(F)$? The answer is NO! In fact, rather, only very few supercuspidal representations are realized in this way. The supercuspidal representations constructed here are called *depth-zero supercuspidal representations*.

4.4. Depth of representations. Let us describe a general picture in the following. We first consider the structure of $\mathrm{GL}_1(F) = F^\times$. As reviewed in Week 2, we have the following isomorphism depending on the choice of a uniformizer ϖ of F^\times :

$$\begin{aligned} F^\times &\cong \langle \varpi \rangle \times \mathcal{O}^\times \\ &\cong \langle \varpi \rangle \times \mathbb{F}_q^\times \times (1 + \mathfrak{p}). \end{aligned}$$

The last part $1 + \mathfrak{p}$ is a profinite group having a descending filtration

$$(1 + \mathfrak{p}) \supset (1 + \mathfrak{p}^2) \supset \cdots \supset \{1\}.$$

This filtration gives a fundamental system of neighborhood. In particular, if a character χ of F^\times is smooth, then it must be trivial on $1 + \mathfrak{p}^r$ for some $r \in \mathbb{Z}_{\geq 0}$. By noting this, we introduce a numerical invariant for smooth characters of F^\times as follows:

Definition 4.5. Suppose that $r \in \mathbb{Z}_{\geq 0}$ is the smallest integer such that χ is trivial on $1 + \mathfrak{p}^{r+1}$ but nontrivial on $1 + \mathfrak{p}^r$. (For convenience, we put $1 + \mathfrak{p}^0 := \mathcal{O}^\times$.) We call this number r the *depth* (or *level*) of χ .

The idea is to generalize this argument to $\mathrm{GL}_2(F)$ (or even more general p -adic reductive groups). The following descending filtration gives a fundamental system of neighborhood of 1:

$$\begin{pmatrix} \mathcal{O} & \mathcal{O} \\ \mathcal{O} & \mathcal{O} \end{pmatrix} \supset \begin{pmatrix} 1 + \mathfrak{p} & \mathfrak{p} \\ \mathfrak{p} & 1 + \mathfrak{p} \end{pmatrix} \supset \begin{pmatrix} 1 + \mathfrak{p}^2 & \mathfrak{p}^2 \\ \mathfrak{p}^2 & 1 + \mathfrak{p}^2 \end{pmatrix} \supset \cdots \supset \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Let us put $K_r := \begin{pmatrix} 1 + \mathfrak{p}^r & \mathfrak{p}^r \\ \mathfrak{p}^r & 1 + \mathfrak{p}^r \end{pmatrix}$ ($K_0 := \mathrm{GL}_2(\mathcal{O})$). The difference between GL_1 and GL_2 is that, for GL_2 , an irreducible representation (π, V) may not be trivial on K_r for any $r \in \mathbb{Z}_{\geq 0}$. (This is because V could be infinite-dimensional; in fact, it happens as long as V is not 1-dimensional.) However, it is still true that V^{K_r} is nonzero for some $r \in \mathbb{Z}_{\geq 0}$ due to the definition of smoothness. Therefore, it still makes sense to look at the smallest integer $r \in \mathbb{Z}_{\geq 0}$ satisfying $V^{K_r} = 0$ but $V^{K_{r+1}} \neq 0$.

Then is it reasonable to define the “depth” of an irreducible smooth representation (π, V) to be this number r ? Actually, NOT! The reason is that $\mathrm{GL}_2(\mathcal{O})$ has many interesting/important open compact subgroups other than K_r 's. Let us consider the following descending chain:

$$\begin{pmatrix} \mathcal{O}^\times & \mathcal{O} \\ \mathfrak{p} & \mathcal{O}^\times \end{pmatrix} \supset \begin{pmatrix} 1 + \mathfrak{p} & \mathcal{O} \\ \mathfrak{p} & 1 + \mathfrak{p} \end{pmatrix} \supset \begin{pmatrix} 1 + \mathfrak{p} & \mathfrak{p} \\ \mathfrak{p}^2 & 1 + \mathfrak{p} \end{pmatrix} \supset \begin{pmatrix} 1 + \mathfrak{p}^2 & \mathfrak{p} \\ \mathfrak{p}^2 & 1 + \mathfrak{p}^2 \end{pmatrix} \supset \cdots \supset \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

We write I_0 for the most left-hand side subgroup and call it the *Iwahori* subgroup of $\mathrm{GL}_2(F)$. We give a half-integer numbering on each subgroup of this filtration as follows:

$$I_r := \begin{cases} \begin{pmatrix} 1 + \mathfrak{p}^s & \mathfrak{p}^s \\ \mathfrak{p}^{s+1} & 1 + \mathfrak{p}^s \end{pmatrix} & \text{if } r = s, \\ \begin{pmatrix} 1 + \mathfrak{p}^{s+1} & \mathfrak{p}^s \\ \mathfrak{p}^{s+1} & 1 + \mathfrak{p}^{s+1} \end{pmatrix} & \text{if } r = s + \frac{1}{2}, \end{cases}$$

where $s \in \mathbb{Z}_{\geq 0}$. Why this way of numbering is reasonable? The point is the second step subgroup is the pro- p -radical of I_0 , i.e., maximal normal pro- p subgroup of I_0 . In this sense, it is an analogue of K_1 for K_0 . However, even if we raise the level of each entry of I_0 , we do not get this second step subgroup; what we get is the third step subgroup. So it is fair to call the third step one “ I_1 ” and the second step one “ $I_{\frac{1}{2}}$ ”.

Definition 4.6. Let (π, V) be an irreducible smooth representation of $\mathrm{GL}_2(F)$. Suppose that $r \in \mathbb{Z}_{\geq 0}$ is the smallest integer such that $V^{P^r} = 0$ but $V^{P^{r+1}} \neq 0$ for $P = K$ or $P = I$. We call the number r the *depth* of the representation (π, V) .

The notion of depth can be generalized for any irreducible smooth representation of any p -adic reductive group; it was introduced by Moy–Prasad [MP94, MP96]. The subgroups K_0 and I_0 are generalized to so-called *parahoric subgroups* of p -adic reductive groups, which can be classified by *Bruhat–Tits theory* [BT72, BT84]. Roughly speaking, Bruhat–Tits classified maximal open compact subgroups of a p -adic reductive group by introducing a geometric object equipped with an action of the p -adic group, which is called the *Bruhat–Tits building*. In the papers of Moy–Prasad [MP94, MP96], they introduced a descending filtration to each such maximal open compact subgroup, which is called the *Moy–Prasad filtration*. (The above filtrations $\{K_r\}_r$ and $\{I_r\}_r$ are nothing but the Moy–Prasad filtrations for K_0 and I_0 .) Then, Moy–Prasad defined the notion of a depth using the all possible Moy–Prasad filtrations.

In general, the depth is known to be a non-negative rational number. Moreover, its possible denominator is determined by the given p -adic reductive group. For example, in the case of GL_2 , the denominator can be only 1 or 2.

4.5. Simple supercuspidal representations. Now let us go back to how to think about supercuspidal representations of $\mathrm{GL}_2(F)$. The representation $\pi_{\tilde{\kappa}}$ constructed above has a non-zero K_1 -fixed vector, thus its depth is zero. In fact, it is known that for any positive half-integer r , there exists an irreducible supercuspidal representation of $\mathrm{GL}_2(F)$ of depth r .

In contrast to the case of finite fields, classifying all positive-depth irreducible supercuspidal representations of $\mathrm{GL}_2(F)$ is not easy nor elementary at all. It’s doable, but based on very subtle and deep analysis of the group structure of $\mathrm{GL}_2(F)$. Because I’m not going to go into its details in this course, here let’s just cite Chapter 4 of Bushnell–Henniart’s book [BH06]. The construction/classification given there can be thought of as a special case of Bushnell–Kutzko’s *type theory* for GL_n [BK93].

But, instead, I just would like to explain how the minimal positive depth (i.e., depth $\frac{1}{2}$) supercuspidal representations can be constructed because it’s fairly easy.

Recall that we have

$$I_0 = \begin{pmatrix} \mathcal{O}^\times & \mathcal{O} \\ \mathfrak{p} & \mathcal{O}^\times \end{pmatrix} \supset I_{\frac{1}{2}} = \begin{pmatrix} 1 + \mathfrak{p} & \mathcal{O} \\ \mathfrak{p} & 1 + \mathfrak{p} \end{pmatrix} \supset I_1 = \begin{pmatrix} 1 + \mathfrak{p} & \mathfrak{p} \\ \mathfrak{p}^2 & 1 + \mathfrak{p} \end{pmatrix}.$$

Note that the quotient $I_{\frac{1}{2}}/I_1$ is isomorphic to the abelian group $\mathbb{F}_q^{\oplus 2}$ by looking at the mod- \mathfrak{p} reduction of $(1, 2)$ and $(2, 1)$ entries:

$$I_{\frac{1}{2}}/I_1 \xrightarrow{\cong} \mathbb{F}_q^{\oplus 2} : \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto (\bar{b}, \overline{c\varpi^{-1}}).$$

We choose a nontrivial additive character $\psi: \mathbb{F}_p \rightarrow \mathbb{C}^\times$ and define a smooth character of $I_{\frac{1}{2}}$ by

$$I_{\frac{1}{2}} \twoheadrightarrow I_{\frac{1}{2}}/I_1 \cong \mathbb{F}_q^{\oplus 2} \xrightarrow{\text{sum}} \mathbb{F}_q \xrightarrow{\psi} \mathbb{C}^\times : \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \psi(\bar{b} + \overline{c\varpi^{-1}}).$$

By abuse of notation, let us again write ψ for this character.

Basically we want to get an irreducible supercuspidal representation of $\text{GL}_2(F)$ by applying the compact induction to this representation of $I_{\frac{1}{2}}$. As in the depth-zero case, we extend ψ to a bit bigger subgroup. The intermediate group we need is the following:

$$\text{GL}_2(F) \supset Z(F) \cdot I_{\frac{1}{2}} \cdot \langle \varphi \rangle \supset I_{\frac{1}{2}}, \quad \varphi = \begin{pmatrix} 0 & 1 \\ \varpi & 0 \end{pmatrix}.$$

We choose any extension $\tilde{\psi}$ of ψ to this subgroup and put

$$\pi_{\tilde{\psi}} := \text{c-Ind}_{Z(F)I_{\frac{1}{2}}\langle\varphi\rangle}^{\text{GL}_2(F)} \tilde{\psi}.$$

Fact 4.7. *The representation $\pi_{\tilde{\psi}}$ is an irreducible supercuspidal representation of $\text{GL}_2(F)$ of depth $\frac{1}{2}$. Conversely, any such representation is of the form $\pi_{\tilde{\psi}}$.*

The representations obtained in this way are called *simple supercuspidal representations* and have discovered firstly by Gross–Reeder [GR10].

Exercise 4.8. Prove that the normalizer group of I_0 in $\text{GL}_2(F)$ is given by $Z(F) \cdot I_0 \cdot \langle \varphi \rangle$.

Exercise 4.9. Describe all the possible extensions of ψ from $I_{\frac{1}{2}}$ to $Z(F)I_{\frac{1}{2}}\langle\varphi\rangle$.

5. WEEK 5: REPRESENTATION OF WEIL GROUPS

For this week's discussion, we follow [BH06, Chapter 7],

5.1. Representations absolute Galois groups. Let us start with investigating continuous representations of profinite groups on \mathbb{C} -vector spaces.

Proposition 5.1. *Let G be a profinite group. Let (π, V) be a continuous representation of G on a finite-dimensional \mathbb{C} -vector space V . Then the image of $\pi: G \rightarrow \text{Aut}_{\mathbb{C}}(V)$ is finite.*

Proof. By choosing a \mathbb{C} -basis of V , we identify $\text{Aut}_{\mathbb{C}}(V)$ with $\text{GL}_n(\mathbb{C})$, where n is the dimension of V . For any positive real number ε , we define an open subset U_ε of $M_n(\mathbb{C})$ by

$$U_\varepsilon := \{g = (g_{ij}) \in M_n(\mathbb{C}) \mid |g_{ij}| < \varepsilon\}.$$

If we choose ε to be sufficiently small, then $I_n + U_\varepsilon$ is a subset of $\text{GL}_n(\mathbb{C})$. Moreover, it is an open neighborhood of I_n in $\text{GL}_n(\mathbb{C})$. Let us write K_ε for this open subset of $\text{GL}_n(\mathbb{C})$. Since π is continuous, the preimage $\pi^{-1}(K_\varepsilon)$ is also an open neighborhood of $1 \in G$.

Recall that G is profinite, hence it has a fundamental system of open neighborhood of $1 \in G$ consisting of open compact subgroups. In fact, even stronger, we can choose such a system so that each subgroup is normal (see Exercise below). Thus let us take an open normal compact subgroup K of G such that $K \subset \pi^{-1}(K_\varepsilon)$, or equivalently, $\pi(K) \subset K_\varepsilon$. Then $\pi(K)$ is a subgroup contained in K_ε .

However, as long as ε is sufficiently small, K_ε does not contain any nontrivial subgroup. Indeed, for the sake of contradiction, let us assume that K_ε contains a nontrivial subgroup K' and choose $k \in K' \setminus \{I_n\}$. Let $\alpha_1, \dots, \alpha_n$ be the generalized eigenvalues of k . Then $\alpha_1^r, \dots, \alpha_n^r$ are the generalized eigenvalues of k^r for any $r \in \mathbb{Z}$. Note that all the eigenvalues of any element of K_ε must be sufficiently close to 1. Since K' is a subgroup contained in K_ε , k^r belongs to K_ε for any $r \in \mathbb{Z}$. Hence $\alpha_1^r, \dots, \alpha_n^r$ are sufficiently close to 1 for any $r \in \mathbb{Z}$. This can happen only when $\alpha_1 = \dots = \alpha_n = 1$. In other words, k must be a unipotent matrix. However, if k is not equal to I_n , k^r cannot belong to K_ε for sufficiently large $r \in \mathbb{Z}_{>0}$. (This can be easily seen by, e.g., taking the Jordan normal form of k).

Hence we obtained that $\pi(K)$ must be $\{I_n\}$. Thus π factors through the quotient G/K , which is a finite group, hence $\pi(G)$ is finite. \square

Exercise 5.2. For any profinite group, prove that there exists a fundamental system of open neighborhoods of 1 consisting of open normal subgroups.

Our fundamental interest lies in understanding the absolute Galois groups of a non-archimedean local field (or even a global field). We approach to this by investigating representations of the absolute Galois group. Since the absolute Galois group is a topological group equipped with a profinite topology, it is natural to impose some topological constraints on the representations. However, the above proposition is saying that “as long as we consider continuous representations on \mathbb{C} -vector spaces, we cannot construct any interesting (nontrivial) examples beyond those coming from finite Galois groups”.

Then, what should be the next candidates for the coefficients and the class of representations to be studied? In our context, one natural idea is to consider continuous representations on $\overline{\mathbb{Q}_\ell}$ -vector spaces. This is because, for example, theory of étale cohomology provides machinery to systematically construct such a class of representations. Also, indeed, there exist plenty of continuous representations with infinite images.

Example 5.3. Let $G = \mathbb{Z}_\ell$. If we define a 2-dimensional representation $\rho: \mathbb{Z}_\ell \rightarrow \text{GL}_2(\overline{\mathbb{Q}_\ell})$ by $x \mapsto \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}$, then π is obviously continuous and has infinite image.

Now we consider the case where G is the absolute Galois group $\text{Gal}(F^{\text{sep}}/F)$ of a non-archimedean local field F . So we investigate continuous representations of $\text{Gal}(F^{\text{sep}}/F)$ on finite-dimensional $\overline{\mathbb{Q}}_\ell$ -vector spaces, where ℓ is a prime number. If we let p be the residual characteristic of F , the situation changes according to whether $\ell = p$ or $\ell \neq p$. In fact, the case where $\ell = p$ is much more complicated and difficult. Although the continuous representations of $\text{Gal}(F^{\text{sep}}/F)$ on $\overline{\mathbb{Q}}_p$ -vector spaces (often referred to as p -adic Galois representations) are very important object to be studied, we focus only on the case where $\ell \neq p$ in this course. In the following, when we say “an ℓ -adic representation”, it means a continuous representation on a finite-dimensional $\overline{\mathbb{Q}}_\ell$ where ℓ is not equal to p .

But then we come up with another question: when $\ell \neq p$, how does the situation depend on the choice of ℓ ? Recall that the other side of the Langlands correspondence (globally, automorphic representations; locally, irreducible admissible representations of a p -adic reductive group) does not involve such a choice of a prime number ℓ . In fact, *Grothendieck’s monodromy theorem* provides an answer to this question.

5.2. Galois group vs. Weil group. Let F be a non-archimedean local field with residual characteristic $p > 0$. Recall that we have a natural surjection

$$\Gamma_F := \text{Gal}(F^{\text{sep}}/F) \twoheadrightarrow \text{Gal}(\overline{k}_F/k_F) \cong \hat{\mathbb{Z}}.$$

The kernel of this surjection is referred to as the *inertia subgroup* and denoted by I_F ; this is nothing but the absolute Galois group $\text{Gal}(F^{\text{sep}}/F^{\text{ur}})$ of the maximal unramified extension F^{ur} of F (see Week 2 notes). The *Weil group* W_F , which is a subgroup of Γ_F , is defined to be the preimage of $\langle \text{Frob}_{k_F} \rangle$ under the map $\Gamma_F \twoheadrightarrow \text{Gal}(\overline{k}_F/k_F)$:

$$\begin{array}{ccccccc} 1 & \longrightarrow & I_F & \longrightarrow & \Gamma_F & \longrightarrow & \text{Gal}(\overline{k}_F/k_F) \cong \hat{\mathbb{Z}} \longrightarrow 1 \\ & & \parallel & & \cup & & \cup \\ 1 & \longrightarrow & I_F & \longrightarrow & W_F & \longrightarrow & \langle \text{Frob}_{k_F} \rangle \cong \mathbb{Z} \longrightarrow 1 \end{array}$$

We write v for the map $W_F \rightarrow \mathbb{Z}$.

In the following, we investigate ℓ -adic representation of W_F rather than Γ_F . Note that W_F is dense in Γ_F , hence any ℓ -adic representation ρ of Γ_F is uniquely determined by its restriction to W_F . To be more precise, we let

- $\text{Rep}_{\overline{\mathbb{Q}}_\ell}(\Gamma_F)$ be the set of isomorphism classes of ℓ -adic representations of Γ_F , and
- $\text{Rep}_{\overline{\mathbb{Q}}_\ell}(W_F)$ be the set of isomorphism classes of ℓ -adic representations of W_F .

Then the natural restriction map gives an injection:

$$\text{Rep}_{\overline{\mathbb{Q}}_\ell}(\Gamma_F) \hookrightarrow \text{Rep}_{\overline{\mathbb{Q}}_\ell}(W_F).$$

Thus at least restricting to W_F does not lose information of the original ℓ -adic representations of Γ_F . However, be careful that there are more ℓ -adic representations of W_F than those of Γ_F , i.e., the above map is not surjective.

Then, why do we work with $\text{Rep}_{\overline{\mathbb{Q}}_\ell}(W_F)$ rather than $\text{Rep}_{\overline{\mathbb{Q}}_\ell}(\Gamma_F)$? This is because $\text{Rep}_{\overline{\mathbb{Q}}_\ell}(W_F)$ can be shown to be independent of ℓ in a certain sense as we will see in the following.

Exercise 5.4. Find an example of an ℓ -adic representation of W_F which cannot be extended to an ℓ -adic representation of Γ_F .

5.3. More about Weil groups. Let us explain a bit more about the group I_F . But, before it, recall that a finite extension of non-archimedean local fields E/F is called ramified when its ramification index is greater than 1.

Definition 5.5. Let E/F be a finite extension of non-archimedean local fields with residual characteristic $p > 0$. We say that E/F is *tamely ramified* when its ramification index is prime to p . Otherwise, we say that E/F is *wildly ramified*.

For any integer $n \in \mathbb{Z}_{>0}$ prime to p , there uniquely exists a degree n extension of F^{ur} . Explicitly, this extension is given by adjoining an(y) n -th root of a(ny) uniformizer ϖ of F . This extension is Galois and cyclic; we have an isomorphism

$$\text{Gal}(F^{\text{ur}}(\sqrt[n]{\varpi})/F^{\text{ur}}) \cong \mu_n: \sigma \mapsto \sigma(\sqrt[n]{\varpi})/\sqrt[n]{\varpi},$$

where μ_n denotes the set of n -th roots of unity (in F^{sep}). We put F^{tame} to be the composite of all finite extensions of F^{ur} whose degree is prime to p . Then F^{tame} is a Galois extension of F . In fact, this gives the maximal tamely ramified extension of F . By the above description of the Galois group at each finite level, we have

$$\text{Gal}(F^{\text{tame}}/F^{\text{ur}}) \cong \varprojlim_{(n,p)=1} \mu_n.$$

Note that, by fixing a system of generators of μ_n (i.e., a topological generator of the right-hand side), we also have

$$\varprojlim_{(n,p)=1} \mu_n \cong \varprojlim_{(n,p)=1} \mathbb{Z}/n\mathbb{Z} \cong \prod_{\ell \neq p} \mathbb{Z}_\ell.$$

We let $P_F := \text{Gal}(F^{\text{sep}}/F^{\text{tame}})$ and call it the *wild inertia subgroup*. In fact, P_F is the unique pro- p -Sylow subgroup of I_F . The quotient $I_F/P_F \cong \text{Gal}(F^{\text{tame}}/F^{\text{ur}})$ is often referred to as the *tame inertia group*.

So we obtained the following chains:

$$\begin{array}{ccc} & \Gamma_F & F \\ \Gamma_F/I_F \cong \hat{\mathbb{Z}} & \nabla & \cap \text{ add } n\text{-th roots of } 1 \text{ (} p \nmid n \text{)} \\ & I_F & F^{\text{ur}} \\ I_F/P_F \cong \prod_{\ell \neq p} \mathbb{Z}_\ell & \nabla & \cap \text{ add } n\text{-th roots of } \varpi \text{ (} p \nmid n \text{)} \\ & P_F & F^{\text{tame}} \\ \text{pro-}p \text{ subgroup} & \nabla & \cap \text{ ???} \\ & \{1\} & F^{\text{sep}} \end{array}$$

Since the conjugate action of Γ_F on itself preserves I_F and P_F , Γ_F also acts on the tame inertia group I_F/P_F . As the tame inertia is abelian, this action factors through the quotient Γ_F/I_F . The action of the subgroup W_F/I_F on I_F/P_F is described as follows:

Lemma 5.6. *For any $\tau \in W_F$ and $\sigma \in I_F/P_F$, we have*

$$\tau \sigma \tau^{-1} = \sigma^{q^{-v(\tau)}}.$$

Proof. By the above description of the tame inertia group ($I_F/P_F \cong \text{Gal}(F^{\text{tame}}/F^{\text{ur}}) \cong \varprojlim_{(n,p)=1} \mu_n$), it is enough to check that

$$\tau \sigma \tau^{-1}(\sqrt[n]{\varpi}) = \sigma^{q^{-v(\tau)}}(\sqrt[n]{\varpi}),$$

for each $n \in \mathbb{Z}_0$ prime to p with the above notation.

Since τ preserves ϖ , $\tau(\sqrt[n]{\varpi})$ is again an n -th root of ϖ ; let us write $\tau(\sqrt[n]{\varpi}) = \zeta \cdot \sqrt[n]{\varpi}$ with some $\zeta \in \mu_n$. We also write $\sigma(\sqrt[n]{\varpi}) = \xi \cdot \sqrt[n]{\varpi}$ with $\xi \in \mu_n$. Here, note that F^{ur} contains all roots of unity with prime-to- p order. In particular,

- σ acts on such roots of unity via identity, and
- τ acts on such roots of unity via $q^{-v(\tau)}$ -power (recall that any lift Φ of the *geometric* Frobenius is supposed to have $v(\Phi) = 1$).

Hence we get

$$\tau\sigma\tau^{-1}(\sqrt[n]{\varpi}) = \tau\sigma(\zeta^{-q^{v(\tau)}} \cdot \sqrt[n]{\varpi}) = \tau(\xi \cdot \zeta^{-q^{v(\tau)}} \cdot \sqrt[n]{\varpi}) = \xi^{q^{-v(\tau)}} \cdot \zeta^{-1} \cdot \zeta \cdot \sqrt[n]{\varpi} = \xi^{q^{-v(\tau)}} \cdot \sqrt[n]{\varpi}.$$

On the other hand, we have

$$\sigma^{q^{-v(\tau)}}(\sqrt[n]{\varpi}) = \xi^{q^{-v(\tau)}} \cdot \sqrt[n]{\varpi}.$$

This completes the proof. \square

5.4. Grothendieck's monodromy theorem. Recall that $I_F/P_F \cong \prod_{\ell \neq p} \mathbb{Z}_\ell$. In the following, we fix a prime number ℓ (supposed to be the “ ℓ ” of “ ℓ -adic representations”) and also fix a surjective homomorphism

$$t: I_F/P_F \cong \prod_{\ell \neq p} \mathbb{Z}_\ell \rightarrow \mathbb{Z}_\ell.$$

Note that hence we have $I_F \supset \text{Ker}(t) \supset P_F$.

For any finite-dimensional C -vector space V (where C is any field of characteristic zero) and its nilpotent endomorphism $N \in \text{End}_C(V)$, we put

$$\exp(N) := \sum_{n=0}^{\infty} \frac{N^n}{n!}.$$

Note that, since $N^n = 0$ for sufficiently large n (at least for n greater than $\dim(V)$), this infinite sum is actually a finite sum. Moreover, $\exp(N)$ is a unipotent automorphism of V . Conversely, for any unipotent automorphism $u \in \text{Aut}_C(V)$, we put

$$\log(u) := \sum_{n=1}^{\infty} (-1)^{n-1} \frac{(u-1)^n}{n}.$$

Then this defines a nilpotent endomorphism of V . These operations give the inverse to each other.

Grothendieck's monodromy theorem says that any ℓ -adic representation of W_F is “quasi-unipotent”:

Theorem 5.7 (Grothendieck's monodromy theorem). *Let ρ be an ℓ -adic representation of W_F . Then there exists an open subgroup H of I_F and a unique nilpotent endomorphism $N \in \text{End}_{\overline{\mathbb{Q}}_\ell}(V)$ satisfying*

$$\rho(\sigma) = \exp(t(\sigma) \cdot N)$$

for any element $\sigma \in H$.

Proof. Let us first check the uniqueness. Suppose that we have two pairs (H, N) and (H', N') as in the assertion. Then, for any $\sigma \in H \cap H'$, we have

$$\exp(t(\sigma) \cdot N) = \rho(\sigma) = \exp(t(\sigma) \cdot N').$$

Thus, by applying \log , we get $t(\sigma) \cdot N = t(\sigma) \cdot N'$. Since $H \cap H'$ is open and of finite index in I_F , the restriction of t on $H \cap H'$ cannot be trivial. Hence we can find $\sigma \in H \cap H'$ such that $t(\sigma) \neq 0$, which implies that $N = N'$.

Let us show the existence. In the following, by fixing a $\overline{\mathbb{Q}_\ell}$ -basis of V , we identify $\text{Aut}_{\overline{\mathbb{Q}_\ell}}(V)$ with $\text{GL}_n(\overline{\mathbb{Q}_\ell})$. Hence ρ is regarded as a continuous homomorphism

$$\rho: W_F \rightarrow \text{GL}_n(\overline{\mathbb{Q}_\ell}).$$

For $r \in \mathbb{Z}_{>0}$, we define an open subgroup K_r of $\text{GL}_n(\overline{\mathbb{Q}_\ell})$ by

$$K_r := I_n + \ell^r \cdot M_n(\overline{\mathbb{Z}_\ell}).$$

We define a subgroup J of I_F by $J := \rho^{-1}(K_2) \cap \text{Ker}(t)$. We claim that $\rho(J) = \{I_n\}$. Indeed, note that $\text{Ker}(t)$ is a profinite group whose pro-order is prime-to- ℓ , that is, $\text{Ker}(t)$ does not have a finite quotient whose order is divided by ℓ . As ρ is continuous and I_F is compact (hence so is J), $\rho(J)$ must be a compact subgroup of K_2 . Since K_2/K_3 is isomorphic to $M_n(\overline{\mathbb{Z}_\ell}/\ell\overline{\mathbb{Z}_\ell})$, which is a discrete abelian group of exponent ℓ , the image of $\rho(J)$ in the quotient K_2/K_3 is discrete and compact, hence finite. But then its order must be ℓ -power, thus $\rho(J)$ is necessarily trivial. In other words, $\rho(J)$ is contained in K_3 . By repeating this argument for K_3, K_4 , and so on, eventually, we get $\rho(J) = \bigcap_{r>0} K_r = \{I_n\}$.

$$\begin{array}{ccc} I_F & \xrightarrow{\rho|_{I_F}} & \text{GL}_n(\overline{\mathbb{Q}_\ell}) \\ \nabla & & \cup \\ \text{Ker}(t) & & K_1 \\ \cup & & \nabla \\ J & \longrightarrow & K_2 \\ & \searrow & \nabla \\ & & K_3 \\ & \searrow & \nabla \\ & & \vdots \\ & & K_4 \\ & & \nabla \\ & & \vdots \end{array}$$

Since $J = \rho^{-1}(K_2) \cap \text{Ker}(t)$, we have $(\rho^{-1}(K_2) \cap I_F)/J \subset I_F/\text{Ker}(t)$. Hence the restriction of ρ to $\rho^{-1}(K_2) \cap I_F$ factors through the homomorphism $t: I_F/\text{Ker}(t) \xrightarrow{\cong} \mathbb{Z}_\ell$. We let $\phi: t(\rho^{-1}(K_2) \cap I_F) \rightarrow K_2$ be the induced homomorphism:

$$\begin{array}{ccc} \rho^{-1}(K_2) \cap I_F & \xrightarrow{\rho} & K_2 \\ \downarrow & & \uparrow \phi \\ (\rho^{-1}(K_2) \cap I_F)/J & \xrightarrow[t \cong]{} & t(\rho^{-1}(K_2) \cap I_F) \\ \cap & & \cap \\ I_F/\text{Ker}(t) & \xrightarrow[t \cong]{} & \mathbb{Z}_\ell \end{array}$$

Let $\Phi \in W_F$ be any lift of the geometric Frobenius, i.e., an element such that $v(\Phi) = 1$. By Lemma 5.6, we have $\Phi\sigma\Phi^{-1} = \sigma^{-q}$ for any $\sigma \in I_F/P_F$. Hence, for any $\sigma \in \rho^{-1}(K_2) \cap I_F$, we have

$$\rho(\Phi)\rho(\sigma)\rho(\Phi)^{-1} = \rho(\Phi\sigma\Phi^{-1}) = \rho(\sigma^{-q}) = \rho(\sigma)^{-q}.$$

This equality implies that the set of eigenvalues of $\rho(\sigma)$ is stable under taking the q -power. Note that this only happens when every eigenvalue of $\rho(\sigma)$ is a root of unity. On the other hand, since $\rho(\sigma) \in K_2$, every eigenvalue of $\rho(\sigma)$ belongs to $1 + \ell^2\overline{\mathbb{Z}_\ell}$. A fun fact here is that these imply that any eigenvalue must be 1 (see exercise below; the reason why we are

looking at “ K_2 ” (not “ K_1 ”) is coming from here). Therefore, $\rho(\sigma)$ must be unipotent for any $\sigma \in \rho^{-1}(K_2) \cap I_F$.

Now we note that $\rho^{-1}(K_2)$ is open in W_F by the continuity of ρ , hence $(\rho^{-1}(K_2) \cap I_F)/J$ is also open (thus of finite index) in $I_F/\text{Ker}(t)$. In particular, $t(\rho^{-1}(K_2) \cap I_F)$ cannot be zero. Hence we can choose $\sigma_0 \in \rho^{-1}(K_2) \cap I_F$ such that $t(\sigma_0) \neq 0$. Let us define a nilpotent endomorphism $N \in M_n(\overline{\mathbb{Q}}_\ell)$ by

$$N := t(\sigma_0)^{-1} \cdot \log(\rho(\sigma_0)).$$

Then obviously we have $\exp(t(\sigma_0) \cdot N) = \rho(\sigma_0) = \phi(t(\sigma_0))$. Let us consider an open (hence of finite index) subgroup $A := t(\sigma_0) \cdot \mathbb{Z}_\ell$ of \mathbb{Z}_ℓ . By definition, we have

$$A \subset t(\rho^{-1}(K_2) \cap I_F) \subset \mathbb{Z}_\ell.$$

We claim that, for any $x \in A$,

$$\exp(x \cdot N) = \phi(x) \in K_2.$$

Indeed, as remarked above, this identity holds for $x = t(\sigma_0)$. Since both \exp and ϕ are multiplicative, then the identity holds for any $x \in t(\sigma_0) \cdot \mathbb{Z} \subset t(\sigma_0) \cdot \mathbb{Z}_\ell = A$. As both \exp and ϕ are continuous, the identity holds for any $x \in t(\sigma_0) \cdot \mathbb{Z}_\ell = A$ (simply because \mathbb{Z} is dense in \mathbb{Z}_ℓ).

We finally put H to be the preimage of A under the map

$$\rho^{-1}(K_2) \cap I_F \twoheadrightarrow (\rho^{-1}(K_2) \cap I_F)/J \xrightarrow{t} t(\rho^{-1}(K_2) \cap I_F).$$

Then H is open in $\rho^{-1}(K_2) \cap I_F$, hence also in I_F . By the observation in the previous paragraph, we have

$$\rho(\sigma) = \phi(t(\sigma)) = \exp(t(\sigma) \cdot N)$$

for any $\sigma \in H$. □

Exercise 5.8. Let $\alpha \in \overline{\mathbb{Q}}_\ell$ be a root of unity such that $\alpha \in 1 + \ell^2 \overline{\mathbb{Z}}_\ell$. Prove that $\alpha = 1$.

5.5. Weil–Deligne representations.

Definition 5.9. Let C be any algebraically closed field of characteristic 0. An n -dimensional *Weil–Deligne representation* of W_F with C -coefficient is a triple (r, V, N) consisting of

- (1) an n -dimensional smooth C -representation (r, V) of W_F ,
- (2) a nilpotent endomorphism $N \in \text{End}_C(V)$ (“monodromy operator”) satisfying

$$r(\sigma) \cdot N \cdot r(\sigma)^{-1} = q^{-v(\sigma)} \cdot N \quad \text{for any } \sigma \in W_F.$$

We can define the notion of a homomorphism (and so on) for Weil–Deligne representations in a natural way. We write WD_C for the set of isomorphism classes of finite-dimensional Weil–Deligne representations with C -coefficients.

Remark 5.10. Recall that the smoothness is equivalent to the continuity with respect to the *discrete* topology of the coefficient field. Thus the choice of C does not matter so much in the above definition. To be more precise, if we have an isomorphism $C \cong C'$ (as abstract fields), then we have $\text{WD}_C \cong \text{WD}_{C'}$.

Exercise 5.11. In fact, any endomorphism $N \in \text{End}_{\mathbb{C}}(V)$ satisfying the condition as in Definition 5.9 (2) is necessarily nilpotent; prove this.

Let (ρ, V) be an ℓ -adic representation of W_F . Let N be the nilpotent endomorphism associated to ρ by Grothendieck's monodromy theorem. We fix a lift $\Phi \in W_F$ of the geometric Frobenius and define a map

$$r: W_F \rightarrow \text{Aut}_{\overline{\mathbb{Q}}_\ell}(V)$$

by

$$r(\Phi^k \cdot \sigma) := \rho(\Phi^k \cdot \sigma) \cdot \exp(t(\sigma) \cdot N)^{-1}$$

for $k \in \mathbb{Z}$ and $\sigma \in I_F$. Then, it is not difficult to see that r is a homomorphism. By the monodromy theorem, r is trivial on an open subgroup of I_F . In other words, (r, V) is a smooth representation of W_F . Furthermore, it can be also checked that (r, V, N) is a Weil–Deligne representation.

Conversely, for any Weil–Deligne representation, we can define an ℓ -adic representation by reversing the above procedure.

Theorem 5.12 (“Second form” of Grothendieck's monodromy theorem). *The above association $\rho \mapsto (r, V, N)$ gives an equivalence between*

- *the category of ℓ -adic representations of W_F , and*
- *the category of finite-dimensional Weil–Deligne representations.*

In particular, we obtain a bijective map

$$\text{WD}: \text{Rep}_{\overline{\mathbb{Q}}_\ell}(W_F) \xrightarrow{1:1} \text{WD}_{\overline{\mathbb{Q}}_\ell}.$$

In fact, it is not difficult to prove that the “converse direction” association $(r, V, N) \mapsto \rho$ gives a well-defined functor and also that it is a fully faithful. So the nontrivial point of the above theorem is that this association can indeed exhausts all ℓ -adic representation; this is nothing but the content of Grothendieck's monodromy theorem. We omit the details of the proof of Theorem 5.12, but it is a routine work as long as we admit Grothendieck's monodromy theorem, which we already proved. See, e.g., [BH06, 32.6].

As mentioned above, the point here is that $\text{WD}_{\overline{\mathbb{Q}}_\ell}$ is essentially independent of ℓ ; for any distinct ℓ' (not equal to p), we have an abstract field isomorphism $\overline{\mathbb{Q}}_\ell \cong \overline{\mathbb{Q}}_{\ell'}$, hence $\text{WD}_{\overline{\mathbb{Q}}_\ell} \cong \text{WD}_{\overline{\mathbb{Q}}_{\ell'}}$. Thus, now we arrived at the following picture.

$$\begin{array}{ccc} \text{Rep}_{\overline{\mathbb{Q}}_\ell}(\Gamma_F) \hookrightarrow & \text{Rep}_{\overline{\mathbb{Q}}_\ell}(W_F) & \xrightarrow[\text{rest}]{1:1} \text{WD}_{\overline{\mathbb{Q}}_\ell} \\ & & \downarrow \scriptstyle 1:1 \quad \overline{\mathbb{Q}}_\ell \cong \overline{\mathbb{Q}}_{\ell'} \\ \text{Rep}_{\overline{\mathbb{Q}}_{\ell'}}(\Gamma_F) \hookrightarrow & \text{Rep}_{\overline{\mathbb{Q}}_{\ell'}}(W_F) & \xrightarrow[\text{rest}]{1:1} \text{WD}_{\overline{\mathbb{Q}}_{\ell'}} \end{array}$$

6. WEEK 6: LOCAL LANGLANDS CORRESPONDENCE FOR GL_n

This week we discuss the statement of the local Langlands correspondence for $\mathrm{GL}(n)$, especially, its characterization.

6.1. Local Langlands correspondence for GL_n . Let F be a non-archimedean local field with residue field $k_F = \mathbb{F}_q$, which is of characteristic p .

Recall that, last week we discussed the notion of Weil–Deligne representation of W_F . In the following, when we talk about a smooth representation of W_F , we always assume that it is finite-dimensional without particularly declaring.

Lemma 6.1. *For any Weil–Deligne representation (r, V, N) , the following are equivalent:*

- (1) *The image of $r(\Phi)$ is semisimple for some lift Φ of the geometric Frobenius.*
- (2) *The image of $r(\Phi)$ is semisimple for any lift Φ of the geometric Frobenius.*
- (3) *The smooth representation (r, V) of W_F is semisimple.*

Proof. Here we omit the proof; see, e.g., [BH06, 32.7]. (Basically the idea is to go back to the proof of monodromy theorem.) \square

Definition 6.2. Let (ρ, V, N) be a Weil–Deligne representation.

- (1) We say that (r, V, N) is *Frobenius-semisimple* if the image of $r(\Phi)$ is semisimple for a lift Φ of the geometric Frobenius.
- (2) We say that (r, V, N) is *semisimple* if it is Frobenius-semisimple and $N = 0$.

Remark 6.3. Note that our terminology is a bit confusing; when a Weil–Deligne representation (r, V, N) is Frobenius-semisimple and N is nonzero, (r, V, N) is not semisimple in our sense, but its underlying smooth representation (r, V) of W_F is semisimple.

We let

- $\Pi(\mathrm{GL}_n)$ be the set of irreducible admissible representations of $\mathrm{GL}_n(F)$, and
- $\mathrm{WD}_{\mathbb{C}, n}^{\mathrm{Frob}}$ be the set of isomorphism classes of 2-dimensional Frobenius-semisimple Weil–Deligne representations.

The local Langlands correspondence for GL_n , which was established by Harris–Taylor and Henniart, asserts that there is a natural bijection between these two sets.

Theorem 6.4 (LLC for GL_n , [HT01, Hen00]). *There exists a unique bijection*

$$\mathrm{LLC}_{\mathrm{GL}_n} : \Pi(\mathrm{GL}_n) \xrightarrow{1:1} \mathrm{WD}_{\mathbb{C}, n}^{\mathrm{Frob}}$$

satisfying the following properties:

- (1) *(compatibility with LCFT) For any $\chi \in \Pi(\mathrm{GL}_1)$, we have*

$$\mathrm{LLC}_{\mathrm{GL}_1}(\chi) = \chi \circ \mathrm{Art}_F^{-1},$$

where $\mathrm{Art}_F : F^\times \cong W_F^{\mathrm{ab}}$ denotes the local Artin map of the local class field theory.

- (2) *(compatibility with character twist) For any $\pi \in \Pi(\mathrm{GL}_n)$ and $\chi \in \Pi(\mathrm{GL}_1)$, we have*

$$\mathrm{LLC}_{\mathrm{GL}_n}(\pi \otimes (\chi \circ \det)) = \mathrm{LLC}_{\mathrm{GL}_n}(\pi) \otimes \mathrm{LLC}_{\mathrm{GL}_1}(\chi).$$

- (3) *(compatibility with central characters) For any $\pi \in \Pi(\mathrm{GL}_n)$ with central character $\omega_\pi \in \Pi(\mathrm{GL}_1)$, we have*

$$\mathrm{LLC}_{\mathrm{GL}_1}(\omega_\pi) = \det \circ \mathrm{LLC}_{\mathrm{GL}_n}(\pi).$$

- (4) *(compatibility with duality) For any $\pi \in \Pi(\mathrm{GL}_n)$, we have*

$$\mathrm{LLC}_{\mathrm{GL}_n}(\pi^\vee) = \mathrm{LLC}_{\mathrm{GL}_n}(\pi)^\vee.$$

(5) (preservation of local factors) For any $\pi_1 \in \Pi(\mathrm{GL}_{n_1})$ and $\pi_2 \in \Pi(\mathrm{GL}_{n_2})$, we have

$$L(s, \pi_1 \times \pi_2) = L(s, \mathrm{LLC}_{\mathrm{GL}_{n_1}}(\pi_1) \otimes \mathrm{LLC}_{\mathrm{GL}_{n_2}}(\pi_2)),$$

$$\varepsilon(s, \pi_1 \times \pi_2) = \varepsilon(s, \mathrm{LLC}_{\mathrm{GL}_{n_1}}(\pi_1) \otimes \mathrm{LLC}_{\mathrm{GL}_{n_2}}(\pi_2)).$$

Here, the left-hand sides are the automorphic local factors of Jacquet–Piatetski-Shapiro–Shalika [JPSS83] and the right-hand sides are the Galois-theoretic local factors of Deligne–Langlands [Del73].

Note that although the properties (1)–(4) are quite important, they do not determine the map $\mathrm{LLC}_{\mathrm{GL}_n}$ uniquely at all. For the unique characterization, the property (5) is really essential.

6.2. Example: the case of GL_2 . Before we discuss the property (5) of Theorem 6.4, we consider the case of GL_2 .

Recall that irreducible admissible representations of $\mathrm{GL}_2(F)$ are classified as follows (Week 4):

- (1) Irreducible principal series representations. The representation $\mathrm{Ind}_{B(F)}^{G(F)} \chi_1 \boxtimes \chi_2$ is irreducible if and only if $\chi_1 \chi_2^{-1} \neq \mathbb{1}, |\cdot|^{-2}$.
- (2) Character twists of Steinberg/trivial representations. If $\chi_1 \chi_2^{-1} = \mathbb{1}$, hence $\chi_1 = \chi_2 = \chi_0$ for some smooth character $\chi_0: F^\times \rightarrow \mathbb{C}^\times$, then $\mathrm{Ind}_{B(F)}^{G(F)} \chi_1 \boxtimes \chi_2 \cong \mathrm{Ind}_{B(F)}^{G(F)} \mathbb{1} \otimes (\chi_0 \circ \det)$ has two irreducible subquotients $\chi_0 \circ \det$ and $\mathrm{St}_{\mathrm{GL}_2} \otimes (\chi_0 \circ \det)$.
- (3) Irreducible supercuspidal representations. The representations which are not of the above two types are called supercuspidal representations.
 - Depth-zero supercuspidal representations.
 - Simple supercuspidal representations (depth $\frac{1}{2}$).
 - Deeper-depth supercuspidal representations.

Let us also classify 2-dimensional semisimple Weil–Deligne representations. Let (r, V, N) be such a representation.

When $N = 0$, we only have two possibilities; (r, V) is an irreducible 2-dimensional smooth representation of W_F or the sum of two smooth 1-dimensional representations (characters) of W_F . Here, we do not talk about how to further classify irreducible 2-dimensional smooth representations of W_F .

We consider the case where $N \neq 0$. In this case, we may choose a basis of V to regard $V \cong \mathbb{C}^{\oplus 2}$ such that the matrix representation of N is given by $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. Then, for any $\sigma \in W_F$, we can check that the form of $r(\sigma)$ is very limited as follows.

Exercise 6.5. Prove that the conditions

- $r(\sigma)$ is semisimple,
- $r(\sigma) \cdot N \cdot r(\sigma) = q^{-v(\sigma)} \cdot N$

implies that

$$r(\sigma) = \begin{pmatrix} z \cdot q^{-\frac{v(\sigma)}{2}} & 0 \\ 0 & z \cdot q^{\frac{v(\sigma)}{2}} \end{pmatrix}$$

for some $z \in \mathbb{C}^\times$.

Let $|\cdot|: W_F \rightarrow \mathbb{C}^\times$ be the absolute value character, i.e., $|\sigma| := q^{-v(\sigma)}$. Then the above observation implies that we must have

$$r = (\chi \otimes |\cdot|^{\frac{1}{2}}) \oplus (\chi \otimes |\cdot|^{-\frac{1}{2}}),$$

where χ is a smooth character of W_F . In other words, if we define a 2-dimensional Frobenius-semisimple Weil–Deligne representation “Sp(2)” by

$$\mathrm{Sp}(2) := (|\cdot|^{1/2} \oplus |\cdot|^{-1/2}, \mathbb{C}^{\oplus 2}, N),$$

then we have $(r, V, N) = \mathrm{Sp}(2) \otimes \chi$.

Now the local Langlands correspondence for GL_2 is stated more precisely as follows:

- (1) An irreducible principal series representation $\mathrm{Ind}_{B(F)}^{G(F)} \chi_1 \boxtimes \chi_2$ corresponds to

$$(\chi_1 \otimes |\cdot|^{1/2} \oplus \chi_2 \otimes |\cdot|^{-1/2}, \mathbb{C}^{\oplus 2}, 0).$$

- (2) A character $\chi \circ \det$ corresponds to

$$(\chi \otimes |\cdot|^{1/2} \oplus \chi \otimes |\cdot|^{-1/2}, \mathbb{C}^{\oplus 2}, 0).$$

- (3) A character twist of the Steinberg representation $\mathrm{St}_{\mathrm{GL}_2} \otimes (\chi \circ \det)$ corresponds to

$$\mathrm{Sp}(2) \otimes \chi = (\chi \otimes |\cdot|^{1/2} \oplus \chi \otimes |\cdot|^{-1/2}, \mathbb{C}^{\oplus 2}, N).$$

- (4) An irreducible supercuspidal representation corresponds to

$$(r, V, 0),$$

where (r, V) is a 2-dimensional irreducible smooth representation of W_F .

6.3. Idea of the characterization of LLC for GL_n . The fundamental philosophy in number theory is:

we should be able to attach a ζ -function or L -function to any number-theoretic object.

Because this is just a slogan, the meaning of “ ζ/L -function” or “number-theoretic object” are not clear. Please just keep in mind that the most basic example is the Riemann ζ -function $\zeta(s)$. So a ζ/L -function in general is something expected to satisfy various nice properties similarly to $\zeta(s)$, e.g., meromorphic continuation to the whole plane \mathbb{C} , functional equation, Euler product decomposition into local factors, and so on. If you have studied theory of modular forms, please remember that we can associate the L -function to any nice modular form and that they indeed satisfy such properties.

Recall that, in Week 1, we looked at an example of the global Langlands correspondence, which relates a modular form (say f) and an elliptic curve (say E). In fact, the mysterious connection between them explained there can be stated in a cleaner way by appealing to their L -functions, i.e., $L(s, f) = L(s, E)$. The point here is that the relation between f and E can be uniquely characterized by this equation (this is a consequence of so-called “strong multiplicity one theorem” on the automorphic side and “Chebotarev density theorem” on the Galois side).

So the idea of formulating the local Langlands correspondence for GL_n is to introduce a local version of L -functions (called “local L -factors”) for irreducible admissible representations of $\mathrm{GL}_n(F)$ and Weil–Deligne representations and then characterize the correspondence using them. However, in fact, using only local L -factors is not enough for the unique characterization. We additionally need “local ε -factors” and also their further variants for “pairs” of representations.

Theory of local factors is too deep to be explained within just one week, so please let me first declare that my explanation below is very naive.

6.4. Local L -factors and ε -factors. Let us start with the following well-known elementary lemma.

Lemma 6.6. *Let G be a group and (ρ, V) be a representation of G . For any normal subgroup H of G , the subspace V^H of H -fixed vectors is G -stable, i.e., a G -subrepresentation. Moreover, the action of G on V^H factors through G/H .*

Proof. Let $v \in V^H$. Our task is to show that, for any $g \in G$, $\rho(g)(v)$ again belongs to V^H . For any $h \in H$, we have

$$\rho(h)(\rho(g)(v)) = \rho(g)(\rho(g^{-1}hg)(v)) = \rho(g)(v),$$

hence $\rho(g)(v)$ is fixed by $\rho(h)$ (in the second equality, we used that H is normal in G). The second assertion is obvious. \square

We first define the local L -factor of a smooth representation of W_F . Recall that the inertia subgroup is a normal subgroup of W_F such that W_F/I_F is isomorphic to the subgroup of $\text{Gal}(\bar{k}_F/k_F)$ which is generated by the geometric Frobenius element Frob_{k_F} (inverse to $x \mapsto x^q$). We fix a lift $\Phi \in W_F$ of Frob_{k_F} .

Definition 6.7. Let (r, V) be a semisimple smooth representation of W_F . We define a complex function $L(s, r)$ on $s \in \mathbb{C}$ by

$$L(s, r) := \det(1 - r(\Phi) \cdot q^{-s} \mid V^{I_F})^{-1}.$$

We call $L(s, r)$ the *local L -factor* of (r, V) .

Remark 6.8. Note that $W_F/I_F \cong \mathbb{Z}$, hence any its semisimple representation decomposes into the sum of 1-dimensional characters of W_F/I_F . We say that such a character (i.e., a character of W_F trivial on I_F) an *unramified character*. If we write $V^{I_F} = \bigoplus_{i=1}^r \chi_i$, where each χ_i is an unramified character of W_F , then we get

$$L(s, r) = \prod_{i=1}^r (1 - \chi_i(\Phi) \cdot q^{-s})^{-1}.$$

Example 6.9. Let us give two extremal examples of local L -factors.

- (1) If (r, V) is the trivial representation of W_F , then $L(s, r) = (1 - q^{-s})^{-1}$. Note that, when $q = p$, this is nothing but the local factor of the Riemann ζ -function

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_p (1 - p^{-s})^{-1}.$$

- (2) If (r, V) is an irreducible smooth representation of W_F , then V^{I_F} must be either 0 or V since it is a W_F -subrepresentation of V . If $V^{I_F} = V$, then it means that (r, V) is a 1-dimensional unramified character. Especially, if the dimension of (r, V) is greater than 1, its L -factor is always trivial ($L(s, r) = 1$).

These examples shows that the local L -factor only knows the unramified part of the given representation. From global perspective, this is enough because any “nice” ℓ -adic Galois representation of a global field is unramified at almost all places and uniquely determined by its behavior there by Chebotarev density theorem. However, from local perspective, the local L -factor is not enough for the unique characterization.

We next define the local L -factor for a Weil–Deligne representation.

Lemma 6.10. *Let (r, V, N) be a Frobenius-semisimple Weil–Deligne representation. Let $V^{N=0}$ denote $\text{Ker}(N: V \rightarrow V)$. Then $V^{N=0}$ is stable under the W_F -action, hence is a semisimple smooth representation of W_F .*

Proof. Suppose $v \in V^{N=0}$. Our task is to show that, for any $\sigma \in W_F$, $r(\sigma)(v)$ again belongs to $V^{N=0}$. We have

$$N(r(\sigma)(v)) = r(\sigma)(r(\sigma)^{-1} \cdot N \cdot r(\sigma)(v)) = r(\sigma)(q \cdot N(v)) = r(\sigma)(0) = 0.$$

□

Definition 6.11. Let (ρ, V, N) be a Frobenius-semisimple Weil–Deligne representation. We define a complex function $L(s, (r, V, N))$ on $s \in \mathbb{C}$ by

$$L(s, (r, V, N)) := L(s, V^{N=0}).$$

We call $L(s, (r, V, N))$ the *local L -factor* of (r, V, N) .

Exercise 6.12. Compute $L(s, \text{Sp}(2))$.

We just give a brief comment on “local ε -factors”. Recall that the (completed) Riemann ζ -function $\hat{\zeta}(s)$ satisfies the functional equation $\hat{\zeta}(s) = \hat{\zeta}(1-s)$. Then, should we expect that such a symmetric equation can be satisfied in general by any L -function associated to a sufficiently nice global object? In fact, it’s not literally so in general, but it is expected that the functional equation holds after adding a correction term called the ε -factor. The typical form of the functional equation for a global object X is like

$$L(s, X) = \varepsilon(s, X) \cdot L(1-s, X^\vee).$$

It is expected that $\varepsilon(s, X)$ also decomposes into the product of local factors, and those local factors are called local ε -factors⁵.

This is just a philosophical explanation of the role of local ε -factors. In our context (i.e., smooth representations of W_F and also Weil–Deligne representations), there are axiomatic properties of the local ε -factor $\varepsilon(s, \rho)$, which is a complex function on $s \in \mathbb{C}$. It is proved that the function $\varepsilon(s, \rho)$ always exists and is uniquely characterized by those properties. In the case of smooth characters, its definition was given by Tate (so-called “Tate’s thesis”). In the general case, it is as follows (see, [BH06, Section 29]).

Theorem 6.13. *For any semisimple smooth representation r of W_F , there uniquely exists a complex function $\varepsilon(s, r) \in \mathbb{C}[q^{\pm s}]^\times$ satisfying the following properties:*

- (1) *If r is 1-dimensional, then $\varepsilon(s, r)$ coincides with Tate’s one.*
- (2) *For any two semisimple smooth representations r_1 and r_2 of W_F , we have $\varepsilon(s, r_1 \oplus r_2) = \varepsilon(s, r_1) \varepsilon(s, r_2)$.*
- (3) *For any finite separable extensions $E \supset K \supset F$ and a semisimple smooth representation r of W_E , we have*

$$\frac{\varepsilon(s, \text{Ind}_{W_E}^{W_K} r)}{\varepsilon(s, r)} = \frac{\varepsilon(s, \text{Ind}_{W_E}^{W_K} \mathbb{1}_{W_E})^{\dim r}}{\varepsilon(s, \mathbb{1}_{W_E})^{\dim r}}.$$

Once we define the local ε -factor for semisimple smooth representations of W_F in this way, we can also extend it to any Frobenius-semisimple Weil–Deligne representations; see [BH06, Section 31] for details.

⁵But note that the local L -factors and the local ε -factors are NOT expected to satisfy the local functional equation. For example, when (ρ, V) is a ramified irreducible representation, the L -factors $L(s, \rho)$ and $L(s, \rho^\vee)$ are trivial, but $\varepsilon(s, \rho)$ is nontrivial and knows how “deep” the ramification of ρ is.

Remark 6.14. For a smooth representation ρ of W_F , its local ε -factor is defined by choosing a nontrivial additive character ψ_F of F . For this reason, it is usually denoted by $\varepsilon(s, \rho, \psi_F)$, but here we omit it from the notation. Note that, in (3) of the above theorem, we choose such a character to be $\psi \circ \text{Tr}_{E/F}$ for any finite separable extension E/F by fixing one ψ_F . (The same is true for Weil–Deligne representations.)

Exercise 6.15. Let (r, V) be a semisimple smooth representation of W_E for a finite separable extension E/F . Prove that $L(s, \text{Ind}_{W_E}^{W_F} r) = L(s, r)$.

So far, we have only talked about the local factors on the Galois side. In fact, there is also a parallel picture established on the automorphic side. It was initiated by Tate in the case of GL_1 (the above-mentioned local factors for 1-dimensional characters of W_F are nothing but the “transfer” of Tate’s factors on the automorphic side via local class field theory) and then generalized to GL_n by Godement–Jacquet.

In this course, I do not explain anything about its definition; actually, it is not easy at all even to state the definition of the local factors on the automorphic side. In some sense, this difficulty of providing a definition and the consequences derived from it are in a “trade-off” relationship. On the Galois side, it is quite easy to define the local L -factor. But it is typically so nontrivial to show that those factors indeed satisfy nice properties, especially, global properties such as meromorphic continuation, functional equation, etc. On the automorphic side, it is already a highly nontrivial task to give its definition. But, once the definition is given, we can prove a lot about its properties by appealing to the well-established general theory of automorphic representations.

6.5. Local L -factors and ε -factors for pairs. I finally also give some comments about pairs.

As mentioned above, the local ε -factor enables us to get more information of the given irreducible admissible representation of $\text{GL}_n(F)$ or Frobenius semisimple Weil–Deligne representation. For example, we can read off the depth (a.k.a., conductor/slope on the Galois side) from the local ε -factor. However, it is still not enough to uniquely determine the given representation. In other words, it really happens that two non-isomorphic representations π_1 and π_2 satisfy $L(s, \pi_1) = L(s, \pi_2)$ and $\varepsilon(s, \pi_1) = \varepsilon(s, \pi_2)$.

The idea is to consider “pairs”. Let us first look at the Galois side. Suppose that an irreducible smooth representation (r, V) of W_F whose dimension is greater than 1 is given. Then it is impossible to recover r from $L(s, r)$ because $L(s, r) = 1$ as explained above. However, what will happen if we consider $L(s, r \otimes r')$ for “all” semisimple smooth representations r' of W_F ? By definition of the local L -factor, $L(s, r)$ has a pole at $s = 0$ if and only if r contains the trivial representation of W_F . Hence, $L(s, r \otimes r')$ contains a pole at $s = 0$ if and only if $r \otimes r'$ contains the trivial representation. Note that

$$\text{Hom}_{W_F}(\mathbb{1}, r \otimes r') = \text{Hom}_{W_F}(r^\vee, r').$$

In particular, when r' is irreducible, we see that $L(s, r \otimes r')$ has a pole at $s = 0$ if and only if r^\vee is isomorphic to r' . Therefore, when two irreducible smooth representations r_1 and r_2 of W_F are given, we can distinguish them by looking at the poles of $L(s, r_1 \otimes r)$ and $L(s, r_2 \otimes r)$.

Note that, so that this idea works, we need the notion of “the tensor product”. On the Galois side, for a given semisimple representations r_1 and r_2 of W_F whose dimensions are n_1 and n_2 , we can construct their tensor product representation $r_1 \otimes r_2$, whose dimension is $n_1 n_2$. Thus what we need on the automorphic side is a way of associating an irreducible

smooth representation “ $\pi_1 \otimes \pi_2$ ” of $\mathrm{GL}_{n_1 n_2}(F)$ to any pair of irreducible admissible representations π_1 of $\mathrm{GL}_{n_1}(F)$ and π_2 of $\mathrm{GL}_{n_2}(F)$. Such an a-priori-hypothetical object is called the *Rankin–Selberg product* of π_1 and π_2 . In fact, the Rankin–Selberg product can only make sense after we prove the local Langlands correspondence for GL_n . (Such an operation on the automorphic side which can be defined by appealing to the local Langlands correspondence is in general referred to as the *Langlands functoriality*). However, the point is that it is possible to establish the definition of $L(s, \pi_1 \otimes \pi_2)$ and $\varepsilon(s, \pi_1 \otimes \pi_2)$ without defining $\pi_1 \otimes \pi_2$; they are called *Rankin–Selberg local factors* of Jacquet–Piatetski-Shapiro–Shalika [JPSS83].

REFERENCES

- [Art13] J. Arthur, *The endoscopic classification of representations: Orthogonal and symplectic groups*, American Mathematical Society Colloquium Publications, vol. 61, American Mathematical Society, Providence, RI, 2013.
- [BH06] C. J. Bushnell and G. Henniart, *The local Langlands conjecture for $GL(2)$* , Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 335, Springer-Verlag, Berlin, 2006.
- [BK93] C. J. Bushnell and P. C. Kutzko, *The admissible dual of $GL(N)$ via compact open subgroups*, Annals of Mathematics Studies, vol. 129, Princeton University Press, Princeton, NJ, 1993.
- [BT72] F. Bruhat and J. Tits, *Groupes réductifs sur un corps local*, Inst. Hautes Études Sci. Publ. Math. (1972), no. 41, 5–251.
- [BT84] ———, *Groupes réductifs sur un corps local. II. Schémas en groupes. Existence d’une donnée radicielle valuée*, Inst. Hautes Études Sci. Publ. Math. (1984), no. 60, 197–376.
- [Del73] P. Deligne, *Les constantes des équations fonctionnelles des fonctions L* , Modular functions of one variable, II (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), 1973, pp. 501–597. Lecture Notes in Math., Vol. 349.
- [DL76] P. Deligne and G. Lusztig, *Representations of reductive groups over finite fields*, Ann. of Math. (2) **103** (1976), no. 1, 103–161.
- [DR09] S. DeBacker and M. Reeder, *Depth-zero supercuspidal L -packets and their stability*, Ann. of Math. (2) **169** (2009), no. 3, 795–901.
- [DS05] F. Diamond and J. Shurman, *A first course in modular forms*, Graduate Texts in Mathematics, vol. 228, Springer-Verlag, New York, 2005.
- [GR10] B. H. Gross and M. Reeder, *Arithmetic invariants of discrete Langlands parameters*, Duke Math. J. **154** (2010), no. 3, 431–508.
- [Hen00] G. Henniart, *Une preuve simple des conjectures de Langlands pour $GL(n)$ sur un corps p -adique*, Invent. Math. **139** (2000), no. 2, 439–455.
- [HT01] M. Harris and R. Taylor, *The geometry and cohomology of some simple Shimura varieties*, Annals of Mathematics Studies, vol. 151, Princeton University Press, Princeton, NJ, 2001, With an appendix by Vladimir G. Berkovich.
- [JPSS83] H. Jacquet, I. I. Piatetskii-Shapiro, and J. A. Shalika, *Rankin-Selberg convolutions*, Amer. J. Math. **105** (1983), no. 2, 367–464.
- [Kal19] T. Kaletha, *Regular supercuspidal representations*, J. Amer. Math. Soc. **32** (2019), no. 4, 1071–1170.
- [Mok15] C. P. Mok, *Endoscopic classification of representations of quasi-split unitary groups*, Mem. Amer. Math. Soc. **235** (2015), no. 1108, vi+248.
- [MP94] A. Moy and G. Prasad, *Unrefined minimal K -types for p -adic groups*, Invent. Math. **116** (1994), no. 1-3, 393–408.
- [MP96] ———, *Jacquet functors and unrefined minimal K -types*, Comment. Math. Helv. **71** (1996), no. 1, 98–121.
- [Oi24] M. Oi, *Representation theory of finite groups of Lie type*, lecture note, available at <https://masaooi.github.io/DL.pdf>, 2024.
- [Ser73] J.-P. Serre, *A course in arithmetic*, Graduate Texts in Mathematics, vol. No. 7, Springer-Verlag, New York-Heidelberg, 1973, Translated from the French.
- [Ser79] ———, *Local fields*, Graduate Texts in Mathematics, vol. 67, Springer-Verlag, New York-Berlin, 1979, Translated from the French by Marvin Jay Greenberg.
- [Wei74] A. Weil, *Basic number theory*, third ed., Die Grundlehren der mathematischen Wissenschaften, vol. Band 144, Springer-Verlag, New York-Berlin, 1974.

DEPARTMENT OF MATHEMATICS, NATIONAL TAIWAN UNIVERSITY, ASTRONOMY MATHEMATICS BUILDING
5F, No. 1, SEC. 4, ROOSEVELT RD., TAIPEI 10617, TAIWAN
Email address: masaooi@ntu.edu.tw